

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example** 

DOB 12/2/1972 Gender: Unknown

**Patient Identifiers:** 01234567890ABCD, 012345

**Visit Number (FIN):** 01234567890ABCD **Collection Date:** 00/00/0000 00:00

## Anti-sp100 and anti-gp210 Antibodies, IgG

ARUP test code 3002482

Anti-gp210 Antibody, IgG

2.0 Units (Ref Interval: 0.0-24.9)

REFERENCE INTERVAL: Anti-gp210 Antibody, IgG

20.0 Units or less.......Regative 20.1-24.9 Units......Equivocal 25.0 Units or greater....Positive

GP210 IgG antibodies can be detected in patients with primary biliary cholangitis (PBC) and may be of diagnostic relevance in a subset of patients with PBC who are negative for anti-mitochondrial antibodies (AMA). These antibodies have a relatively low sensitivity with excellent specificity for PBC. A negative result does not rule out PBC.

Anti-sp100 Antibody, IgG

4.5 Units

(Ref Interval: 0.0-24.9)

REFERENCE INTERVAL: Anti-sp100 Antibody, IgG

20.0 Units or less.......Negative 20.1-24.9 Units......Equivocal 25.0 Units or greater....Positive

SP100 IgG antibodies can be detected in patients with primary biliary cholangitis (PBC) and may be of diagnostic relevance in a subset of patients with PBC who are negative for anti-mitochondrial antibodies (AMA). These antibodies have a relatively low sensitivity with excellent specificity for PBC. A negative result does not rule out PBC.

H=High, L=Low, \*=Abnormal, C=Critical



VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Anti-gp210 Antibody, IgG	24-046-135395	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Anti-sp100 Antibody, IgG	24-046-135395	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Patient: Patient, Example ARUP Accession: 24-046-135395 Patient Identifiers: 01234567890ABCD, 012345 Visit Number (FIN): 01234567890ABCD Page 2 of 2 | Printed: 2/26/2024 8:42:44 AM

4848