

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 3/6/1948
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

IDH1 and IDH2 Mutation Analysis Exon 4, Formalin-Fixed, Paraffin-Embedded (FFPE) Tissue

ARUP test code 3004267

IDH1 and IDH2 Mutation Results

Not Detected

A mutation was not detected in the IDH1 (codon R132) or IDH2 (codons R140 or R172) genes. This does not exclude the possibility of a mutation below the limit of detection for this assay.

This result has been reviewed and approved by [REDACTED]

BACKGROUND INFORMATION: IDH1 and IDH2 Mutation Results

CHARACTERISTICS: This test is designed to detect mutations in exon 4 of the IDH1 and IDH2 genes at "hotspots" R132 of IDH1 and R140 and R172 of IDH2 that are frequently present in gliomas and in a subset of cases of acute myeloid leukemia. IDH1/2 mutations in gliomas are generally associated with a better prognosis. In acute myeloid leukemia, the prognostic significance of IDH1 mutations is context dependent. IDH1 mutations appear to be associated with worse outcome in patients without FLT3-ITD mutations (see J Clin Oncol 2010. 28:3636 and Blood 2010. 116:2779). In acute myeloid leukemia patients with IDH2 abnormalities, IDH2 R140 mutations appear to be associated with better outcome while IDH2 R172 mutations appear associated with worse outcome (see Blood 2011. 118:409). The FDA has approved ivosidenib as a targeted therapy for acute myeloid leukemia with an IDH1 mutation, and enasidenib for AML with an IDH2 mutation. Clinical trials may be available.

METHODOLOGY: DNA is isolated from FFPE tissue, blood, or bone marrow. The DNA is amplified for IDH1 and IDH2 covering exon 4 of both genes including the important residues R132 (IDH1), R140 (IDH2) and R172 (IDH2). Sanger sequencing is then performed to detect mutations. Only mutations in R132 (IDH1), R140 and R172 (IDH2) are reported.

LIMITATIONS: Mutations in other locations within the IDH1 and IDH2 genes or in other genes will not be detected. The limit of detection for this test is 20 percent mutant allele. Results of this test must always be interpreted within the clinical context and with other relevant data, and should not be used alone for a diagnosis of malignancy. This test is not intended to detect minimal residual disease.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

H=High, L=Low, *=Abnormal, C=Critical

IDH1-2 FFPE Source

FFPE Tissue

Block ID

23ON-104 A1

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
IDH1 and IDH2 Mutation Results	23-353-401005	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
IDH1-2 FFPE Source	23-353-401005	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Block ID	23-353-401005	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Block ID	23-353-401005	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical