

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 5/15/2000  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Circulating Immune Complex, C3 fragments**

ARUP test code 3004431

Circulating Immune Complex, C3 fragments **>40 ug Eq/mL H (Ref Interval: <=15)**

This patient's serum shows reactivity in the circulating immune complex (CIC) inhibitory assay performed in conjunction with the CIC-C3 fragment assay. Reactivity in the inhibitory assay indicates the presence of aggregates of human IgG or other CICs which may be due to: 1) autoantibodies or 2) heat-inactivated or otherwise compromised serum specimens. The presence of concomitantly occurring CICs cannot be confirmed in this patient.

**INTERPRETIVE INFORMATION: Circulating Immune Complex, C3 fragments**

Many autoimmune disorders, chronic infections and malignancies are associated with circulating immune complexes. Quantitation of immune complexes assists in staging immunologic disorders. Detection of circulating immune complexes is not essential to any specific diagnosis. Circulating immune complexes may be found without any evident pathology and positive results do not necessarily implicate immune complex-related disease process. Values between 15 and 20 ug Eq/mL are considered equivocal for Circulating Immune Complex, C3 fragments assay. Repeat- testing using a new specimen is recommended, if clinically indicated.

**VERIFIED/REPORTED DATES**

Procedure	Accession	Collected	Received	Verified/Reported
Circulating Immune Complex, C3 fragments	22-136-118454	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at: