

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: Unknown
Gender: Unknown
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Autoimmune Encephalopathy/Dementia Panel, Serum

ARUP test code 3006201

Neuronal Antibody (Amphiphysin)

Negative (Ref Interval: Negative)

INTERPRETIVE INFORMATION: Amphiphysin Antibody, IgG

Amphiphysin antibody is present in about 5 percent of patients with stiff-person syndrome and is found variably in other causes of paraneoplastic neurological syndrome (PNS). Amphiphysin antibody is mainly associated with small-cell lung cancer and breast tumors.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Purkinje Cell/Neuronal Nuclear IgG Scrn

None Detected (Ref Interval: None Detected)

ANNA-1, ANNA-2, PCCA-1 or PCCA-Tr(DNER) antibodies not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: Purkinje Cell/Neuronal Nuclear IgG Scrn

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

NMDA Receptor Ab IgG CBA-IFA, Serum

<1:10 (Ref Interval: <1:10)

Antibodies to NMDA were not detected, no additional testing to follow.

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: NMDA Receptor Ab IgG CBA-IFA, Serum

NMDA receptor antibody is found in a subset of patients with autoimmune limbic encephalitis and may occur with or without associated tumor. Decreasing antibody levels may be associated with therapeutic response. In addition, positive results have been reported in patients with non-autoimmune phenotypes. A negative test result does not rule out a diagnosis of autoimmune limbic encephalitis. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings. Serum testing should be paired with CSF testing for improved diagnostic sensitivity.

This indirect fluorescent antibody assay utilizes full-length GluN1 transfected cell lines for the detection and semiquantification of NMDA receptor IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

CASPR2 Ab IgG CBA-IFA Screen, Serum

<1:10 (Ref Interval: <1:10)

CASPR2 Antibody, IgG is not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: CASPR2 Ab IgG CBA-IFA Screen, Serum

Contactin-associated protein-2 (CASPR2) IgG antibody may occur as part of the voltage-gated potassium channel (VGKC) complex antibodies.

The presence of CASPR2 IgG antibody is associated with a wide spectrum of clinical manifestations, including acquired neuromyotonia, limbic encephalitis, painful neuropathy, and Morvan syndrome. Tumors such as thymoma, small cell lung cancer, and other rarer tumors may occur. The full-spectrum of clinical disorders and tumors associated with the CASPR2 IgG antibody continues to be defined. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes CASPR2 transfected cell lines for the detection and semiquantification of the CASPR2 IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

LGI1 Ab IgG CBA-IFA Screen, Serum

<1:10 (Ref Interval: <1:10)

LGI1 Antibody, IgG is not detected. No further testing will be performed.

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: LGI1 Ab IgG CBA-IFA Screen, Serum

Leucine-rich, glioma-inactivated 1 protein (LGI1) IgG antibody may occur as part of the voltage-gated potassium channel (VGKC) complex antibodies.

The presence of LGI1 IgG antibody is mainly associated with limbic encephalitis, hyponatremia, and myoclonic movements. LGI1 IgG antibody is rarely associated with tumors but may occur infrequently in Morvan syndrome, neuromyotonia, and idiopathic epilepsy. The full-spectrum of clinical disorders associated with the LGI1 IgG antibody continues to be defined. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes LGI1 transfected cell lines for the detection and semiquantification of the LGI1 IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

CV2 Ab IgG CBA-IFA Screen, Serum

<1:100 (Ref Interval: <1:100)

CV2 Antibody, IgG is not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: CV2 Ab IgG CBA-IFA Screen, Serum

CV2 antibodies aid in discriminating between chronic paraneoplastic neurological disorder (PND) and other inflammatory disorders of the nervous system. Anti-CV2 is associated with small-cell lung cancer and thymoma. A negative test result does not rule out a diagnosis of autoimmune neurologic disease. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes CV2 transfected cell lines for the detection and semiquantification of the CV2 IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

AMPA Receptor Ab IgG CBA-IFA Scrn, Serum

<1:10 (Ref Interval: <1:10)

AMPA Antibody, IgG is not detected. No further testing will be performed.

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: AMPA Receptor Ab IgG CBA-IFA Scrn, Serum

Alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) antibody is found in a subset of patients with autoimmune limbic encephalitis and may occur with or without associated tumor. Decreasing antibody levels may be associated with therapeutic response. A negative test result does not rule out a diagnosis of autoimmune encephalitis. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes AMPAR transfected cell lines for the detection and semiquantification of AMPAR IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

GABA-BR Ab IgG CBA-IFA Scrn, Ser

<1:10 (Ref Interval: <1:10)

GABA-BR Antibody, IgG is not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: GABA-BR Ab IgG CBA-IFA Scrn, Ser

Gamma-amino butyric acid receptor, type B (GABA-BR) antibody is found in a subset of patients with autoimmune epilepsy and other autoimmune neurologic phenotypes; it may occur with or without associated tumor. Decreasing antibody levels may be associated with therapeutic response. A negative test result does not rule out a diagnosis of autoimmune neurologic disease. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes GABA-BR transfected cell lines for the detection and semiquantification of GABA-BR IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

SOX1 Antibody, IgG by Immunoblot, Serum

Negative (Ref Interval: Negative)

INTERPRETIVE INFORMATION: SOX1 Antibody, IgG by Immunoblot, Serum

SOX1 antibody is detected in patients with Lambert-Eaton myasthenic syndrome (LEMS) and in patients with paraneoplastic cerebellar degeneration (PCD), paraneoplastic and nonparaneoplastic neuropathy. SOX1 antibody is associated with small cell lung cancer. A negative test result does not rule out a diagnosis of LEMS or other causes of paraneoplastic neurological syndrome.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

DPPX Ab IgG CBA-IFA Screen, Serum

<1:10 (Ref Interval: <1:10)

H=High, L=Low, *=Abnormal, C=Critical

DPPX Antibody, IgG is not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: DPPX Ab IgG CBA-IFA Screen, Serum

DPPX antibody is found in a subset of patients with autoimmune encephalitis, and is often associated with prodromal gastrointestinal symptoms and unintentional weight loss. It may occur with or without associated tumor. Decreasing antibody levels may be associated with therapeutic response. A negative test result does not rule out a diagnosis of autoimmune neurologic disease. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes DPPX transfected cells for the detection and semiquantification of the DPPX IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

IgLON5 Ab IgG CBA-IFA Screen, Serum

<1:10 (Ref Interval: <1:10)

IgLON5 Antibody, IgG is not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: IgLON5 Ab IgG CBA-IFA Screen, Serum

IgLON Family Member 5 (IgLON5) antibody is found in a subset of patients with autoimmune encephalitis or other autoimmune neurologic/neurodegenerative disorders and may occur with or without associated tumor. A negative test result does not rule out a diagnosis of an autoimmune neurologic disorder. Interpretation of any antineural antibody test requires clinical correlation.

This indirect fluorescent antibody assay utilizes IgLON5 transfected cell lines for detection and semi-quantification of IgLON5 IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

mGluR1 Ab IgG CBA-IFA Screen, Serum

<1:10 (Ref Interval: <1:10)

mGluR1 Antibody, IgG is not detected. No further testing will be performed.

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: mGluR1 Ab IgG CBA-IFA Screen,
Serum

Metabotropic glutamate receptor 1 (mGluR1) antibody is found in a subset of patients with autoimmune cerebellar ataxia or autoimmune encephalitis and may occur with or without associated tumor. A negative test result does not rule out a diagnosis of autoimmune cerebellar ataxia or limbic encephalitis. Interpretation of any antineural antibody test requires clinical correlation.

This indirect fluorescent antibody assay utilizes mGluR1 transfected cell lines for detection and semi-quantification of mGluR1 IgG antibody.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

Glutamic Acid Decarboxylase Antibody

<5.0 IU/mL (Ref Interval: 0.0-5.0)

INTERPRETIVE INFORMATION: Glutamic Acid Decarboxylase Antibody

A value greater than 5.0 IU/mL is considered positive for Glutamic Acid Decarboxylase Antibody (GAD Ab). This assay is intended for the semi-quantitative determination of the GAD Ab in human serum. Results should be interpreted within the context of clinical symptoms.

H=High, L=Low, *=Abnormal, C=Critical

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Neuronal Antibody (Amphiphysin)	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Purkinje Cell/Neuronal Nuclear IgG Scrn	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
NMDA Receptor Ab IgG CBA-IFA, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
CASPR2 Ab IgG CBA-IFA Screen, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
LGI1 Ab IgG CBA-IFA Screen, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
CV2 Ab IgG CBA-IFA Screen, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
AMPA Receptor Ab IgG CBA-IFA Sern, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GABA-BR Ab IgG CBA-IFA Scrn, Ser	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SOX1 Antibody, IgG by Immunoblot, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
DPPX Ab IgG CBA-IFA Screen, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
IgLON5 Ab IgG CBA-IFA Screen, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
mGluR1 Ab IgG CBA-IFA Screen, Serum	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Glutamic Acid Decarboxylase Antibody	24-057-100498	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 24-057-100498
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 7 of 7 | Printed: 2/26/2024 11:13:06 AM
4848