

Patient: Client:	ARUP Test Code: 300637
------------------	------------------------

DOB: Age: Sex:

Patient Identifiers:

Collection Date: 10/09/2023
Physician:

Received in lab: 10/10/2023
Visit Number (FIN):

Completion Date: 10/11/2023

Patient Results*

Collection Date 10/9/2023 3:50 AM

THC-COOH Present

Interpretive Comments

INTERPRETIVE INFORMATION: Drug Detection Panel, Umbilical Cord Tissue, Qualitative

Methodology: Qualitative Liquid Chromatography/Tandem Mass Spectrometry

Detection of drugs in umbilical cord tissue is intended to reflect maternal drug use during approximately the last trimester of a full-term pregnancy. The pattern and frequency of drug(s) used by the mother cannot be determined by this test. A negative result does not exclude the poss bility that a mother used drugs during pregnancy. Detection of drugs in umbilical cord tissue depends on extent of maternal drug use, as well as drug stability, unique characteristics of drug deposition in umbilical cord tissue, and the performance of the analytical method. Drugs administered during labor and delivery may be detected. Detection of drugs in umbilical cord tissue does not insinuate impairment and may not affect outcomes for the infant. Interpretive questions should be directed to the laboratory.

Refer to the test directory for additional umbilical cord testing options.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

INTERPRETIVE INFORMATION: Marijuana Metabolite, Umbilical Cord Tissue, Qualitative

Methodology: Qualitative Liquid Chromatography-Tandem Mass Spectrometry

This test is designed to detect and document exposure that occurred during approximately the last trimester of a full term pregnancy, to a common cannabis (marijuana) metabolite. Alternative testing is available to detect other drug exposures. The pattern and frequency of drug(s) used by the mother cannot be determined by this test. A negative result does not exclude the poss bility that a mother used drugs during pregnancy. Detection of drugs in umbilical cord tissue depends on extent of maternal drug use, as well as drug stability, unique characteristics of drug deposition in umbilical cord tissue, and the performance of the analytical method. Drugs administered during labor and delivery may be detected. Detection of drugs in umbilical cord tissue does not insinuate impairment and may not affect outcomes for the infant. Interpretive questions should be directed to the laboratory.









Patient:

^{*} For details, see report on following page(s).

Patient: | Date of Birth: | Sex: | Physician: Patient Identifiers: | Visit Number (FIN):

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

For further information, visit: arupconsult.com/content/newborn-drug-testing









Patient:

Patient: | Date of Birth: | Sex: | Physician:
Patient Identifiers: | Visit Number (FIN):

·	, ,		
Drug Analyte	Result	Cutoff	Notes
OPIOIDS, UMBILICAL CORD			
Buprenorphine	Not Detected	1 ng/g	eg, Buprenex; Subutex; Suboxone (with naloxone)
Norbuprenorphine	Not Detected	0.5 ng/g	buprenorphine metabolite
Codeine	Not Detected	0.5 ng/g	
Morphine	Not Detected	0.5 ng/g	also a metabolite of Codeine
6-Acetylmorphine	Not Detected	1 ng/g	heroin metabolite; morphine is expected
Hydrocodone	Not Detected	0.5 ng/g	also a minor metabolite of codeine; possible impurity of oxycodone
Dihydrocodeine	Not Detected	1 ng/g	hydrocodone metabolite
Norhydrocodone	Not Detected	1 ng/g	hydrocodone metabolite
Hydromorphone	Not Detected	0.5 ng/g	eg, Dilaudid; also a metabolite of hydrocodone, and a minor metabolite of morphine
Fentanyl	Not Detected	0.5 ng/g	
Meperidine	Not Detected	2 ng/g	eg, Demerol
Methadone	Not Detected	2 ng/g	
EDDP	Not Detected	1 ng/g	methadone metabolite
Oxycodone	Not Detected	0.5 ng/g	eg, Oxycontin
Noroxycodone	Not Detected	1 ng/g	oxycodone metabolite
Oxymorphone	Not Detected	0.5 ng/g	eg, Opana; also a metabolite of oxycodone
Noroxymorphone	Not Detected	0.5 ng/g	oxymorphone metabolite; also potential metabolite of oxycodone
Naloxone	Not Detected	1 ng/g	opioid antagonist
Propoxyphene	Not Detected	1 ng/g	eg, Darvon
Tapentadol	Not Detected	2 ng/g	eg, Nucynta
Tramadol	Not Detected	2 ng/g	eg, Ultram; ConZip
N-desmethyltramadol	Not Detected	2 ng/g	tramadol metabolite
O-desmethyltramadol	Not Detected	2 ng/g	tramadol metabolite
STIMULANTS, UMBILICAL CORD			
Amphetamine	Not Detected	5 ng/g	eg, Vyvanse; Adderall; also a metabolite of methamphetamine; d- and l-isomers are not distinguished by this test
Methamphetamine	Not Detected	5 ng/g	d- and l- isomers are not distinguished by this test
MDMA - Ecstasy	Not Detected	5 ng/g	
Cocaine	N (D ()	1 ng/g	
Cocame	Not Detected	1 119/9	
Benzoylecgonine	Not Detected Not Detected	0.5 ng/g	cocaine metabolite









Patient:

Patient: Patient Identifiers:	Date of Birth: Visit Number (FIN):	Sex:	Physician:
Drug Analyte	Result	Cutoff	Notes
Cocaethylene	Not Detected	1 ng/g	cocaine metabolite formed in the presence of ethanol
Phentermine	Not Detected	8 ng/g	
SEDATIVES-HYPNOTICS UMBILICAL CORD	S,		
Alprazolam	Not Detected	0.5 ng/g	eg, Xanax
Alpha-OH-Alprazolam	Not Detected	0.5 ng/g	alprazolam metabolite
Butalbital	Not Detected	25 ng/g	eg, Bupap (with acetaminophen); Fioricet (with acetaminophen and caffeine)
Clonazepam	Not Detected	1 ng/g	eg, Klonopin
7-Aminoclonazepam	Not Detected	1 ng/g	clonazepam metabolite
Lorazepam	Not Detected	5 ng/g	eg, Ativan
Midazolam	Not Detected	1 ng/g	eg, Versed
Alpha-OH-Midazolam	Not Detected	2 ng/g	midazolam metabolite
Diazepam	Not Detected	1 ng/g	eg, Valium
Nordiazepam	Not Detected	1 ng/g	metabolite of chlordiazepoxide (Librium), clorazepate (Tranxene), diazepam, halazepam (Alapryl), prazepam (Centrax) and others
Oxazepam	Not Detected	2 ng/g	eg, Serax; also metabolite of nordiazepam and temazepam
Temazepam	Not Detected	1 ng/g	eg, Restoril; also a metabolite of diazepam
Phenobarbital	Not Detected	75 ng/g	eg, Luminal; Donnatal (with atropine, hyoscyamine and scopolamine)
Zolpidem	Not Detected	0.5 ng/g	eg, Ambien
OTHER, UMBILICAL CO	RD		
Phencyclidine- PCP	Not Detected	1 ng/g	phencyclidine
Gabapentin	Not Detected	10 ng/g	eg, Neurontin or Gralise
ТНС-СООН	Present	0.2 ng/g	11-Nor-9-carboxy-THC, a marijuana metabolite

Additional Information

ABBREVIATIONS

MDMA - 3,4-methylenedioxy-N-methylamphetamine EDDP - 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine

THC-COOH - cannot discriminate between delta-8 and delta-9 forms









Patient: