

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 3/1/1922
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Tissue Transglutaminase Antibody, IgA

ARUP test code 3016860

Tissue Transglutaminase (tTG) Ab, IgA

<1.02 FLU (Ref Interval: 0.00-4.99)

INTERPRETIVE INFORMATION: Tissue Transglutaminase (tTG) Antibody, IgA

Presence of the tissue transglutaminase (tTG) IgA antibody is associated with gluten-sensitive enteropathies such as celiac disease and dermatitis herpetiformis. Individuals with positive results should be confirmed with small intestinal biopsy to establish celiac disease diagnosis. tTG IgA antibody concentrations greater than 50 FLU exhibits higher correlation with results of duodenal biopsies consistent with celiac disease. For antibody concentrations greater than or equal to 5 FLU but less than 10 FLU, additional testing for endomysial (EMA) IgA concentrations may improve the positive predictive value for disease. A decrease in tTG IgA antibody concentration after initiation of a gluten-free diet may indicate a response to therapy.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Tissue Transglutaminase (tTG) Ab, IgA	23-324-112010	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: