

Client: ARUP Example Report Only
500 Chipeta Way
Salt Lake City, UT 84108
UNITED STATES

Physician: Jones, Shalane

Patient: Meas, EncephSerPos

DOB

Sex: Male

Patient Identifiers: 74822

Visit Number (FIN): 75255

Collection Date: 12/11/2025 14:31

Encephalitis Panel With Reflex to Herpes Simplex Virus Types 1 and 2 Glycoprotein G-Specific Antibodies, IgG, Serum

ARUP test code 3017751

HSV Type 1/2 Combined Ab, IgG

5.00 IV H (Ref Interval: <=0.89)

Specimen tested positive for Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG. ARUP test codes 0050292 and 0050294 will be added. Additional charges apply.

HSV Type 1 and Type 2 Glycoprotein G-Specific Antibodies, IgG to follow.

INTERPRETIVE INFORMATION: HSV 1/2 COMBINED Ab SCREEN, IgG

0.89 IV or less.....Not Detected
0.90-1.09 IV.....Indeterminate- Repeat testing in 10-14 days may be helpful.
1.10 IV or greater.....Detected

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

West Nile Virus Ab, IgG, Ser

5.00 IV H (Ref Interval: <=1.29)

INTERPRETIVE INFORMATION: West Nile Virus Ab, IgG by ELISA, Serum

1.29 IV or less Negative - No significant level of west Nile virus IgG antibody detected.
1.30 - 1.49 IV Equivocal - Questionable presence of West Nile virus IgG antibody detected. Repeat testing in 10-14 days may be helpful.
1.50 IV or greater Positive - Presence of IgG antibody to west Nile virus detected, suggestive of current or past infection.

This test is intended to be used as a semi-quantitative means of detecting west Nile virus-specific IgG in serum samples in which there is a clinical suspicion of west Nile virus infection. This test should not be used solely for quantitative purposes, nor should the results be used without correlation to clinical history or other data. Because other members of the Flaviviridae family, such as St.Louis encephalitis virus, show extensive cross-reactivity with west Nile virus, serologic testing specific for these species should be considered.

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Meas, EncephSerPos
ARUP Accession: 25-345-120056
Patient Identifiers: 74822
Visit Number (FIN): 75255
Page 1 of 5 | Printed: 12/11/2025 2:49:55 PM

Seroconversion between acute and convalescent sera is considered strong evidence of current or recent infection. The best evidence for infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

West Nile Virus Ab, IgM, Ser

5.00 IV H (Ref Interval: <=0.89)

INTERPRETIVE INFORMATION: West Nile Virus Ab, IgM by ELISA, Serum

- 0.89 IV or less Negative - No significant level of west Nile virus IgM antibody detected.
- 0.90-1.10 IV Equivocal - Questionable presence of west Nile virus IgM antibody detected. Repeat testing in 10-14 days may be helpful.
- 1.11 IV or greater ... Positive - Presence of IgM antibody to West Nile virus detected, suggestive of current or recent infection.

This test is intended to be used as a semi-quantitative means of detecting west Nile virus-specific IgM in serum samples in which there is a clinical suspicion of west Nile virus infection. This test should not be used solely for quantitative purposes, nor should the results be used without correlation to clinical history or other data. Because other members of the Flaviviridae family, such as St. Louis encephalitis virus, show extensive cross-reactivity with west Nile virus, serologic testing specific for these species should be considered.

Seroconversion between acute and convalescent sera is considered strong evidence of current or recent infection. The best evidence for infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Mumps Virus Antibody, IgG

13.0 AU/mL

INTERPRETIVE INFORMATION: Mumps Ab, IgG by CIA

- 8.9 AU/mL or less Negative - No significant level of detectable IgG mumps virus antibody
- 9.0-10.9 AU/mL Equivocal - Repeat testing in 10-14 days may be helpful
- 11.0 AU/mL or greater: Positive - IgG antibody to mumps virus detected, which may indicate a current or past exposure/immunization to mumps virus.

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Mumps Virus Antibody, IgM

12.00 IV H (Ref Interval: <=0.79)

INTERPRETIVE INFORMATION: Mumps Virus Antibody, IgM

- 0.79 IV or less: Negative - No significant level of detectable IgM antibody to mumps virus.
- 0.80 - 1.20 IV: Equivocal - Borderline levels of IgM antibody to mumps virus. Repeat

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Meas, EncephSerPos
ARUP Accession: 25-345-120056
Patient Identifiers: 74822
Visit Number (FIN): 75255
Page 2 of 5 | Printed: 12/11/2025 2:49:55 PM

1.21 IV or greater: testing in 10-14 days may be helpful. Positive - Presence of IgM antibody to mumps virus detected, which may indicate a current or recent infection. However, low levels of IgM antibody may occasionally persist for more than 12 months post-infection or immunization.

Measles, Rubeola, Antibody IgG

18.0 AU/mL

INTERPRETIVE INFORMATION: Measles (Rubeola) Antibody, IgG

13.4 AU/mL or less..... Negative - No significant level of detectable measles (rubeola) IgG antibody.
13.5-16.4 AU/mL Equivocal - Repeat testing in 10-14 days may be helpful.
16.5 AU/mL or greater Positive - IgG antibody to measles (rubeola) detected which may indicate a current or past exposure/immunization to measles (rubeola).

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Measles, Rubeola, Antibody IgM

1:40 * (Ref Interval: <1:10)

INTERPRETIVE INFORMATION: Measles (Rubeola) Antibody, IgM by IFA

Less than 1:10..... Negative - No evidence of recent infection. False negative results are possible if the specimen was collected too soon after exposure. Molecular testing may be helpful.
1:10 or greater Positive - Indicative of recent primary measles infection. False positive results possible.

Varicella-Zoster Virus Ab, IgG

4.00 S/CO H (Ref Interval: <=0.99)

INTERPRETIVE INFORMATION: VZV Ab, IgG

<=0.99 S/CO: Negative - No significant level of detectable varicella-zoster IgG antibody.
>=1.00 S/CO: Positive - IgG antibody to varicella-zoster detected, which may indicate a current or past varicella-zoster infection.

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Varicella-Zoster Virus Antibody, IgM

5.00 ISR H (Ref Interval: <=0.90)

INTERPRETIVE INFORMATION: Varicella-Zoster Virus Antibody, IgM

0.90 ISR or less Negative - No significant level of detectable

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Meas, EncephSerPos
ARUP Accession: 25-345-120056
Patient Identifiers: 74822
Visit Number (FIN): 75255
Page 3 of 5 | Printed: 12/11/2025 2:49:55 PM

	varicella-zoster virus IgM antibody.
0.91-1.09 ISR	Equivocal - Repeat testing in 10-14 days may be helpful.
1.10 ISR or greater	Positive - Significant level of detectable varicella-zoster virus IgM antibody. Indicative of current or recent infection. However, low levels of IgM antibodies may occasionally persist for more than 12 months post-infection or immunization.

Herpes Simplex Virus Type 2 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050294

HSV 2 Glycoprotein G Antibody, IgG

3.00 IV H (Ref Interval: <=0.89)

REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

0.89 IV or less	Negative - No significant level of detectable IgG antibody to HSV type 2 glycoprotein G.
0.90 - 1.09 IV	Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
1.10 IV or greater	Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

False positive results are possible. Consider additional testing for result less than or equal to 3.0 IV for type specific assays.

Herpes Simplex Virus Type 1 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050292

HSV 1 Glycoprotein G Ab, IgG

3.00 IV H (Ref Interval: <=0.89)

REFERENCE INTERVAL: HSV 1 Glycoprotein G Ab, IgG

0.89 IV or less	Negative - No significant level of detectable IgG antibody to HSV type 1 glycoprotein G.
0.90 - 1.09 IV	Equivocal - Questionable presence of IgG antibody to HSV type 1 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
1.10 IV or greater ...	Positive - IgG antibody to HSV type 1 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

False positive results are possible. Consider additional testing

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Meas, EncephSerPos
ARUP Accession: 25-345-120056
Patient Identifiers: 74822
Visit Number (FIN): 75255
Page 4 of 5 | Printed: 12/11/2025 2:49:55 PM

for result less than or equal to 3.0 IV for type specific assays.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
HSV Type 1/2 Combined Ab, IgG	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:38:00 PM
West Nile Virus Ab, IgG, Ser	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
West Nile Virus Ab, IgM, Ser	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
Mumps Virus Antibody, IgG	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
Mumps Virus Antibody, IgM	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
HSV 1 Glycoprotein G Ab, IgG	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:38:22 PM	12/11/2025 2:47:00 PM
HSV 2 Glycoprotein G Antibody, IgG	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:38:22 PM	12/11/2025 2:47:00 PM
Measles, Rubeola, Antibody IgG	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
Measles, Rubeola, Antibody IgM	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
Varicella-Zoster Virus Ab, IgG	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM
Varicella-Zoster Virus Antibody, IgM	25-345-120056	12/11/2025 2:31:00 PM	12/11/2025 2:34:19 PM	12/11/2025 2:39:00 PM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Meas, EncephSerPos
ARUP Accession: 25-345-120056
Patient Identifiers: 74822
Visit Number (FIN): 75255
Page 5 of 5 | Printed: 12/11/2025 2:49:55 PM