

**Patient: Patient, Example** 

Unknown

Unknown

01234567890ABCD, 012345

01234567890ABCD

00/00/0000 00:00

DOB

Gender:

**Patient Identifiers:** 

Visit Number (FIN):

**Collection Date:** 

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

#### **Polymyositis Panel** ARUP test code 3018868

Jo-1 (Histidyl-tRNA Synthetase) Ab, IgG	0 AU/mL INTERPRETIVE INFORM	(Ref Interval: 0-40) ATION: Jo-1 Antibody, IgG	
	29 AU/mL or lessNegative 30-40 AU/mLEquivocal 41 AU/mL or greaterPositive		
	Presence of Jo-1 (antihistidyl transfer RNA [t-RNA] synthetase) antibody is associated with polymyositis and may also be seen in patients with dermatomyositis. Jo-1 antibody is associated with pulmonary involvement (interstitial lung disease). Raynaud phenomenon, arthritis, and mechanic's hands (implicated in antisynthetase syndrome).		
PL-12 (alanyl-tRNA synthetase) Antibody	Negative	(Ref Interval: Negative)	
PL-7 (threonyl-tRNA synthetase) Antibody	Negative	(Ref Interval: Negative)	
EJ (glycyl-tRNA synthetase) Antibody	Negative	(Ref Interval: Negative)	
OJ (isoleucyl-tRNA synthetase) Antibody	Negative	(Ref Interval: Negative)	
SRP (Signal Recognition Particle) Ab	Negative	(Ref Interval: Negative)	
Polymyositis Interpretive Information	See Note		

# H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:



INTERPRETIVE	INFORMATION:	Polymyositis	Panel	2
--------------	--------------	--------------	-------	---

	<pre>myositis, and may be prognosis. MSAs are with rare exceptions be carefully evaluat presentation. Myosit in patients with CTD generally not specif help in identifying either MSAs or MAAs. Antibody Specificity Jo-1 (histidyl-tRNA PL-12 (alanyl-tRNA sy PL-7 (threonyl-tRNA EJ (glycyl-tRNA synt) OJ (isoleucyl-tRNA synt) OJ (isoleucyl-tRNA synt) Ha (tyrosyl-tRNA synt) Ks (asparaginyl-tRNA</pre>	-specific antibodies (MSAs) are specific for useful in establishing diagnosis as well as generally regarded as mutually exclusive ; the occurrence of two or more MSAs should ed in the context of patient's clinical is-associated antibodies (MAAs) may be found including overlap syndromes, and are ic for myositis. The following table will the association of any antibodies found as MSAs MAAs synthetase) Ab, IgG X ynthetase) Antibody X synthetase) Antibody X ion Particle) Ab X thetase) Ab X synthetase) Ab X
	This test was develo determined by ARUP La approved by the US Fo	A synthetase) Ab X ped and its performance characteristics aboratories. It has not been cleared or ood and Drug Administration. This test was certified laboratory and is intended for
Antinuclear Antibody (ANA), HEp-2, IgG	<1:80	(Ref Interval: <1:80)
ANA Interpretive Comment	speckled, nucleolar, Cytoplasmic antibodi discrete/GW body-lik cytoplasmic speckled INTERPRETIVE INFORMA Presence of antinucle of systemic autoimmu lacks diagnostic spe- diseases (cancers, a conditions) and may varying prevalence. confirmation of posi ANA (nuclear reactiv centromere, homogener ANA (cytoplasmic rea reticular/AMA, discr cytoplasmic speckled are reported to endp help guide different specific for individ patterns not reporter out SARD.	TION: ANA Interpretive Comment ear antibodies (ANA) is a hallmark feature ne rheumatic diseases (SARD). However, ANA cificity and is associated with a variety of utoimmune, infectious, and inflammatory also occur in healthy individuals in The lack of diagnostic specificity requires tive ANA by more specific serologic tests. ity) positive patterns reported include ous, nuclear dots, nucleolar, or speckled. ctivity) positive patterns reported include ete/GW body-like, polar/golgi-like, or rods and rings. All positive patterns may ial diagnosis, although they may not be ual antibodies or diseases. Mitotic staining d. Negative results do not necessarily rule
Ha (tyrosyl-tRNA synthetase) Ab		(Ref Interval: Negative) by line immunoassay. No band corresponding y immunoprecipitation.

### H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:



Ks (asparaginyl-tRNA synthetase) Ab	Negative Ks antibody negative by line to 65 kDa observed by immuno	(Ref Interval: Negative) immunoassay. No band corresponding precipitation.
Zo (phenylalanyl-tRNA synthetase) Ab	Negative Zo antibody negative by line to 68 and 58 kDa observed by	(Ref Interval: Negative) immunoassay. No bands corresponding immunoprecipitation.

VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Jo-1 (Histidyl-tRNA Synthetase) Ab, IgG	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
PL-12 (alanyl-tRNA synthetase) Antibody	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
PL-7 (threonyl-tRNA synthetase) Antibody	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
EJ (glycyl-tRNA synthetase) Antibody	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
OJ (isoleucyl-tRNA synthetase) Antibody	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SRP (Signal Recognition Particle) Ab	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Polymyositis Interpretive Information	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Antinuclear Antibody (ANA), HEp-2, IgG	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
ANA Interpretive Comment	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Ha (tyrosyl-tRNA synthetase) Ab	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Ks (asparaginyl-tRNA synthetase) Ab	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Zo (phenylalanyl-tRNA synthetase) Ab	25-022-102248	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

# END OF CHART

## H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: