

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 8/13/1952  
**Sex:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Trypanosoma cruzi Antibody, IgG Panel**

ARUP test code 3020058

T. cruzi IgG, Recombinant Antigen

**9.4 S/CO H (Ref Interval: <=0.8)**

INTERPRETIVE INFORMATION: T. cruzi IgG, Recombinant Antigen

- 0.8 S/Co or less ..... Negative - No significant level of Trypanosoma cruzi IgG antibody detected.
- 0.9 - 1.1 S/Co ..... Equivocal - Questionable presence of Trypanosoma cruzi IgG antibody detected. Repeat testing in 10-14 days may be helpful.
- 1.2 S/Co or greater ..... Positive - IgG antibodies to Trypanosoma cruzi detected, which may suggest current or past infection.

According to the CDC, at least two different serologic tests should be used to make the laboratory diagnosis of chronic Chagas disease, as no single serologic test is sufficiently sensitive and specific. If results between the two assays are discrepant, repeat testing or testing by a third method may be helpful.

This assay should not be used for blood donor screening or associated reentry protocols, or for screening human cell and cellular tissue-based products (HCT/Ps).

T. cruzi IgG, Purified Antigen

**3.7 IV H (Ref Interval: <=1.0)**

INTERPRETIVE INFORMATION: T. cruzi IgG, Purified Antigen

- 1.0 IV or less.....Negative - No significant level of Trypanosoma cruzi IgG antibody detected.
- 1.1 IV.....Equivocal - Questionable presence of Trypanosoma cruzi IgG antibody detected. Repeat testing in 10-14 days may be helpful.
- 1.2 IV or greater...Positive - IgG antibodies to Trypanosoma cruzi detected, which may suggest current or past infection.

According to the CDC, at least two different serologic tests should be used to make the laboratory diagnosis of chronic Chagas disease, as no single serologic test is sufficiently sensitive and specific. If results between the two assays are discrepant, repeat testing or testing by a third method may be helpful.

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at:

**ARUP LABORATORIES | 800-522-2787 | aruplab.com**  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 25-302-155633  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 1 of 2 | Printed: 11/4/2025 1:02:51 PM

This assay should not be used for blood donor screening or associated reentry protocols, or for screening human cell and cellular tissue-based products (HCT/PS).

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
T. cruzi IgG, Recombinant Antigen	25-302-155633	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
T. cruzi IgG, Purified Antigen	25-302-155633	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at:

**ARUP LABORATORIES | 800-522-2787 | aruplab.com**  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 25-302-155633  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 2 of 2 | Printed: 11/4/2025 1:02:51 PM