

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB 2/16/1995
Sex: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

von Willebrand Factor Multimeric Panel

ARUP test code 3020171

Factor VIII, Activity 167 % (Ref Interval: 56-191)

REFERENCE INTERVAL: Factor VIII, Activity

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

von Willebrand Factor, Activity (GPIbM) 138 % (Ref Interval: 51-215)

INTERPRETIVE INFORMATION: von Willebrand Factor Activity, GPIbM

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

von Willebrand Factor, Antigen 103 % (Ref Interval: 52-214)

REFERENCE INTERVAL: von Willebrand Factor, Antigen

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

von Willebrand Factor Multimers

ARUP test code 0092281

von Willebrand Factor Multimers

See Note

von Willebrand factor multimeric analysis shows a normal multimeric distribution. A von Willebrand panel without VWF or factor VIII deficiencies combined with a normal multimeric distribution does not entirely rule out von Willebrand disease since von Willebrand values can fluctuate over time. If clinically indicated, consider repeating the von Willebrand panel. Additional information regarding diagnosis and subtyping of von Willebrand disease is available at www.arupconsult.com.

von Willebrand factor and factor VIII are acute phase reactants. Increased values are seen in pregnancy, with estrogen-containing medications, inflammation, infection, liver disease, malignancy, and with exercise, stress, or trauma, including traumatic venipuncture. Acute phase reactions can mask low baseline values

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 26-023-124190
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 1 of 2 | Printed: 2/5/2026 9:20:22 AM

in patients with von Willebrand disease.

Lower baseline von Willebrand and factor VIII values are found in blood type O individuals. Decreased values for von Willebrand factor and/or factor VIII activity may also be due to improper specimen handling.

INTERPRETIVE INFORMATION: von Willebrand Factor Multimers

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Factor VIII, Activity	26-023-124190	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
von Willebrand Factor, Activity (GPIbM)	26-023-124190	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
von Willebrand Factor, Antigen	26-023-124190	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
von Willebrand Factor Multimers	26-023-124190	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 26-023-124190
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 2 of 2 | Printed: 2/5/2026 9:20:22 AM