

## TEST CHANGE

### Rapid Plasma Reagin (RPR) with Reflex to Titer and FTA-ABS

0050011, RPR FTA

#### Specimen Requirements:

##### Patient Preparation:

Collect: Serum separator tube.

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 1mL serum to an ARUP standard transport tube. (Min: 0.5 mL) Avoid freezing if possible.

Transport Temperature: Refrigerated.

Unacceptable Conditions: Contaminated, grossly hemolyzed, grossly lipemic, plasma, CSF, cord blood, or other body fluids.

##### Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year

Methodology: Semi-Quantitative Particle Agglutination

Performed: Sun-Sat

Reported: 1-3 days

##### Note:

CPT Codes: 86592; if reflexed, add 86593; 86780

New York DOH Approval Status: This test is New York DOH approved.

#### Interpretive Data:

Component	Interpretation
Rapid Plasma Reagin (RPR)	RPR (+) = Reactive RPR (-) = Nonreactive

#### Reference Interval:

Test Number	Components	Reference Interval		
	Rapid Plasma Reagin (RPR)	<del>Nonreactive</del> Non- Reactive		
	Rapid Plasma Reagin (RPR)			
		Component Result	Interpretation	
		Rapid Plasma	RPR (+) =	

		Reagin (RPR)	Reactive RPR (-) = Nonreactive	