

HOTLINE: Effective May 16, 2022

0050164 Fungal Antibodies by Immunodiffusion

FUNG PPT

Methodology: Immunodiffusion

Specimen Required: Collect: Serum separator tube (SST)

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP Standard Transport Tube. (Min: 0.4 mL)

Storage/Transport Temperature: Refrigerated.

Unacceptable Conditions: Other body fluids. Contaminated, hemolyzed, icteric, or lipemic specimens.

Stability (collection to initiation of testing): After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)

Reference Interval:

Effective May 16, 2022

Test Number	Components	Reference Interval
0050171	<i>Aspergillus</i> spp. Antibodies by Immunodiffusion	None detected
0050172	<i>Blastomyces dermatitidis</i> Antibodies by Immunodiffusion	None detected
	<i>Coccidioides</i> Antibodies by Immunodiffusion	Not Detected
0050174	<i>Histoplasma</i> spp. Antibodies by Immunodiffusion	None detected

HOTLINE NOTE: There is a clinically significant charting name change associated with this test.

Change the charting name for component 0050183, *Coccidioides immitis* Abs, Precipitin from *Coccidioides immitis* Abs, Precipitin to *Coccidioides* by Immunodiffusion, Serum.