

0050171

Aspergillus Antibodies by Immunodiffusion

ASPER PPT

Methodology: Immunodiffusion
Performed: Sun-Sat
Reported: 3-6 days

Specimen Required: Collect: Serum separator tube.

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum an ARUP Standard Transport Tube. (Min: 0.15 mL) **Mark specimens plainly as "acute" or "convalescent."**

Storage/Transport Temperature: Refrigerated.

Unacceptable Conditions: **Contaminated, hemolyzed, or severely lipemic specimens.**

Stability (collection to initiation of testing): After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)

Reference Interval:

Not Detected

Interpretive Data:

Refer to Report.

Note: This immunodiffusion test uses pooled mycelial-phase culture filtrates of *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus niger*, and *Aspergillus terreus*.

HOTLINE NOTE: There is a clinically significant charting name change associated with this test.

Change the charting name for component 0050171, Aspergillus spp. Abs, Precipitin from Aspergillus spp. Abs, Precipitin to *Aspergillus* Antibodies by ID.