

TEST CHANGE

Rapid Plasma Reagin (RPR) with Reflex to Titer

0050471, RPRT

Specimen Requirements:

Patient Preparation:

Collect: Serum separator tube.

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP [standard transport tube](#). ~~Standard Transport Tube~~. (Min: 0.54 mL) Avoid freezing if possible.

Transport Temperature: Refrigerated.

Unacceptable Conditions: Plasma, CSF, or other body fluids.

Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative [Particle](#)~~Chareal~~ Agglutination

Performed: Sun-Sat

Reported: Within 24 hours

Note: If RPR is reactive, then a titer will be added. Additional charges apply.

CPT Codes: 86592; if reflexed, add 86593

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Component	Interpretation
Rapid Plasma Reagin (RPR)	RPR (+) = Reactive RPR (-) = Nonreactive

Reference Interval:

Test Number	Components	Reference Interval		
	Rapid Plasma Reagin (RPR)	Non Reactive		
	Rapid Plasma Reagin (RPR)			
		Component Result	Interpretation	
		Rapid Plasma Reagin (RPR)	RPR (+) = Reactive RPR (-) = Nonreactive	