

TEST CHANGE

Extractable Nuclear Antigen Antibodies (Smith/RNP, Smith, SSA 52, SSA 60, and SSB)
0050652, ENA ABS4

Specimen Requirements:

Patient Preparation:

Collect: Serum separator tube.

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP Standard Transport Tube. (Min: 0.3 mL)

Transport Temperature: Refrigerated..

Unacceptable Conditions: Plasma or other body fluids. Bacterially contaminated or severely lipemic specimens.

Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month (avoid repeated freeze/thaw cycles)

Methodology: Quantitative Enzyme-Linked Immunosorbent Assay/Quantitative Multiplex Bead Assay

Performed: Sun-Sat

Reported: 1-~~3~~² days

Note:

CPT Codes: 86235 x5

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Component	Interpretation
Smith/RNP (ENA) Antibody, IgG	19 Units or less Negative 20-39 Units Weak Positive 40-80 Units Moderate Positive 81 Units or greater Strong Positive
Smith (ENA) Antibody, IgG	29 AU/mL or less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive

SSA-52 (Ro52) (ENA) Antibody, IgG	29 AU/mL or Less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive
SSA-60 (Ro60) (ENA) Antibody, IgG	29 AU/mL or Less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive
SSB (La) (ENA) Antibody, IgG	29 AU/mL or less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive

Reference Interval:

Test Number	Components	Reference Interval
	Smith/RNP (ENA) Ab, IgG	19 Units or less
	Smith (ENA) Antibody, IgG	40 AU/mL or less
	SSA-52 (Ro52) (ENA) Antibody, IgG	40 AU/mL or less
	SSA-60 (Ro60) (ENA) Antibody, IgG	40 AU/mL or less
	SSB (La) (ENA) Antibody, IgG	40 AU/mL or less