TEST CHANGE

Extractable Nuclear Antigen Antibodies (SSA 52, SSA 60, and SSB)

0050791, SSA/SSB

Specimen Requirements:	
Patient Preparation:	
Collect:	Serum separator tube or red tube.
Specimen Preparation:	Separate serum from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP standard transport tube. (Min: 0.5~~3~~ mL)
Transport Temperature:	Refrigerated.
Unacceptable Conditions:	Plasma or other body fluids. Contaminated, hemolyzed, or severely lipemic specimens.
Remarks:	
Stability:	After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)
Methodology:	Semi-Quantitative Multiplex Bead Assay
Note:	
CPT Codes:	86235 x3
New York DOH Approval Status:	This test is New York DOH approved.

Interpretive Data:

Component	Interpretation
SSA-52 (Ro52) (ENA) Antibody, IgG	29 AU/mL or Less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive
SSA-60 (Ro60) (ENA) Antibody, IgG	29 AU/mL or Less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive
SSB (La) (ENA) Antibody, IgG	29 AU/mL or less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive

Reference Interval:

Test Number	Components	Reference Interval
	SSA-52 (Ro52) (ENA) Antibody, IgG	40 AU/mL or less
	SSA-60 (Ro60) (ENA) Antibody, IgG	40 AU/mL or less
	SSB (La) (ENA) Antibody, IgG	40 AU/mL or less