

TEST CHANGE

Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG with Reflex to Type 1 and 2 Glycoprotein G-Specific Ab, IgG

0051708, HERPR PAN2

Specimen Requirements:

Patient Preparation:

Collect: Serum separator tube.

Specimen Preparation: Transfer 1 mL serum to an ARUP **standard transport tube** ~~Standard Transport Tube~~. (Min: 0.5 mL) Parallel testing is preferred and convalescent specimens must be received within 30 days from receipt of acute specimens. Mark specimens plainly as "acute" or "convalescent."

Transport Temperature: Refrigerated.

Unacceptable Conditions: Plasma or urine. Contaminated, heat-inactivated, hemolyzed, **icteric**, or severely lipemic specimens.

Remarks:

Stability: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year

Methodology: Semi-Quantitative Chemiluminescent Immunoassay

Performed: Sun-Sat

Reported: 1-2 days

Note: If HSV 1 and/or 2 IgG is 1.10 IV or greater, then HSV 1 gG-Specific IgG and HSV 2 gG-Specific IgG will be added. Additional charges apply.

CPT Codes: 86694; if reflexed, add 86695; 86696

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

False positive results are possible. Consider additional testing for HSV-2, particularly if the result for HSV-2 is **less than or equal to** \leq 3.0 IV.

Component	Interpretation
Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG by Chemiluminescent Immunoassay	0.89 IV or less: Not Detected. 0.90-1.09 IV: Indeterminate. \rightarrow Repeat testing in 10-14 days may

	be helpful. 1.10 IV or greater: Detected.	
--	---	--

Reference Interval:
