

TEST CHANGE

Inhibin B

0070413, INHIBINB

Specimen Requirements:

Patient Preparation: For premenopausal females, collection is preferred during the follicular phase of the menstrual cycle.

Collect: Serum separator tube or plain red.

Specimen Preparation: Transport 0.5 mL serum. (Min: 0.2 mL)

Transport Temperature: Frozen.

Unacceptable Conditions: Room temperature specimens. Grossly hemolyzed specimens. Plasma

Remarks:

Stability: After separation from cells: Ambient: Unacceptable; Refrigerated: 72 hours; Frozen: 1 month

Methodology: Quantitative Enzyme-Linked Immunosorbent Assay (ELISA)

Performed: Wed, Fri

Reported: 1-8 days

Note:

CPT Codes: 83520

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

This test is performed using the ANSH ultra-sensitive Inhibin B ELISA kit. Values obtained with different methodologies or kits cannot be used interchangeably.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Reference Interval:

Effective February 16, 2021

Male	Female
<15 days: 68-373 pg/mL 15 days-6 months:	1 day-12 years: ≤ 1 -182 pg/mL 13-41 years

42-516 pg/mL 7 months-7 years:	(regular cycle, follicular phase):
24-300 pg/mL 8-30 years:	8-223 pg/mL 42-51 years (regular cycle, follicular phase):
10-357 pg/mL	$\leq 1-107$ pg/mL 51-76 years (postmenopausal):
	$\leq 1-11$ pg/mL