

TEST CHANGE

Babesia microti Antibody, IgG by IFA

0093049, BAB IGG

Specimen Requirements:

Patient Preparation:

Collect: Serum Separator Tube (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum to an ARUP ~~standard transport tube~~ **Standard Transport Tube**. (Min: 0.42 mL) Parallel testing is preferred and convalescent specimens must be received within 30 days from receipt of the acute specimens. Mark specimens plainly as "acute" or "convalescent."

Transport Temperature: Refrigerated.

Unacceptable Conditions: Bacterially contaminated, hemolyzed, or lipemic specimens.

Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative Indirect Fluorescent Antibody **(IFA)**

Performed: Mon, Wed, Sat

Reported: 1-5 days

Note:

CPT Codes: 86753

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

~~This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.~~

Reference Interval:

< 1:16 Negative - No significant level of detectable *Babesia* IgG antibody.

1:16 Equivocal - Repeat testing in 10-14 days may be helpful.

> 1:16 Positive - IgG antibody to *Babesia* detected, which may indicate a current or past infection.

Deleted Cells



*A nonprofit enterprise of the University of Utah
and its Department of Pathology*

Effective Date: **May 15, 2023**