

## TEST CHANGE

Allergen, Fungi and Molds, Candida albicans IgG

0097304, CANDIDAIGG

### Specimen Requirements:

#### Patient Preparation:

Collect: Serum separator tube (SST).

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum to an ARUP [standard transport tube](#). ~~Standard Transport Tube~~. (Min: 0. ~~252~~ mL). [Refer to "Allergen Specimen Collection Instructions" at www.aruplab.com/testing/resources/specimen.](#)

Transport Temperature: Refrigerated.

Unacceptable Conditions: Hemolyzed, icteric, or lipemic specimens.

#### Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year

Methodology: Quantitative ImmunoCAP Fluorescent Enzyme Immunoassay

Performed: Sun

Reported: 1-8 days

Note: The units of measure mcg/mL and mgA/L are interchangeable.  
1 mg/L = 1000 mcg/1000 mL

CPT Codes: 86001

New York DOH Approval Status: Specimens from New York clients will be sent out to a New York DOH approved laboratory, if possible.

#### Interpretive Data:

Values less than 2.00 mcg/mL represent absent or undetectable levels of allergen-specific IgG antibody.

~~This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.~~

#### Reference Interval:

Less than 84.01 mcg/mL

