

TEST CHANGE

Fluphenazine

0099906, FLUPHEN

Specimen Requirements:

Patient Preparation: Timing of specimen collection: Pre-dose (trough) draw - At steady state concentration.

Collect: Plain red. Also acceptable: Lavender (K2 or K3EDTA) or pink (K2 EDTA).

Specimen Preparation: Separate serum or plasma from cells within 2 hours of collection. Transfer 1 mL serum or plasma to an ARUP Standard Transport Tube. (Min: 0.5 mL)

Transport Temperature: Refrigerated.

Unacceptable Conditions: Whole blood. Hemolyzed specimens. Gel separator tubes, light blue (citrate), or yellow (SPS or ACD solution).

Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 1 week; Frozen: 1 month (avoid repeated freeze/thaw cycles)

Methodology: Quantitative Liquid Chromatography-Tandem Mass Spectrometry

Performed: Mon, Wed, Fri

Reported: 1-85 days

Note:

CPT Codes: 80342 (Alt code: G0480)

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

The therapeutic range is based on serum pre-dose (trough) draw at steady-state concentration. Adverse effects may include extrapyramidal symptoms, seizures and neuroleptic malignant syndrome.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Reference Interval:

Effective February 16, 2021

Therapeutic Range:	1.0-10.0 ng/mL	
Toxic:	Greater than 15 ng/mL	