

TEST CHANGE

Protein S, Free Antigen with Reflex to Protein S, Total Antigen

2002269, PRS FREE R

Specimen Requirements:

Patient Preparation:

Collect: Lt. blue (sodium citrate). ~~Special~~Refer to Specimen ~~Collection and Handling Hemostasis/Thrombosis Specimens guide~~ located at <https://www.aruplab.com/Specimen-Handling/SpecialSpecimenCollection/Hemostasis-Thrombosis.pdf> for hemostasis/thrombosis specimen handling guidelines.

Specimen Preparation: Transfer 2 mL platelet-poor plasma to an ARUP ~~standard transport tube~~.Standard Transport Tube. (Min: 1 mL)

Transport Temperature: CRITICAL FROZEN. Separate specimens must be submitted when multiple tests are ordered.

Unacceptable Conditions: Serum. EDTA plasma, clotted or hemolyzed specimens.

Remarks:

Stability: Ambient: 4 hours; Refrigerated: Unacceptable; Frozen ~~at -20 Degrees C: 3 months, at -70 Degrees C: 6 months~~

Methodology: Microlatex Particle-Mediated Immunoassay

Performed: Mon-Sat

Reported: 1-4 days

Note: If low Protein S Free Antigen is detected, then Protein S, Total Antigen, will be added. Additional charges apply.

CPT Codes: 85306; if reflexed, add 85305

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Refer to report

Reference Interval:

1-89 days: 15-55%
90-179 days: 35-92%
180-364 days: 45-115%

1-5 years: 62-120%

6-9 years: 62-130%

10-17 years: 60-140%

18 years and older Male: 74-147%

18 years and older Female: 55-123%
