

HOTLINE: Effective February 1, 2021

2006495

Phosphatidylserine Antibodies, IgG and IgM

PHOSSER GM

**Specimen Required:** Collect: Serum separator tube

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum to an ARUP Standard Transport Tube. (Min: 0.3 mL)

Storage/Transport Temperature: Refrigerated

Unacceptable Conditions: Heat-inactivated, contaminated, grossly icteric, grossly hemolyzed, or severely lipemic specimens

Stability (collection to initiation of testing): After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month

**Reference Interval:**

Effective February 1, 2021

Test Number	Components	Reference Interval
	Phosphatidylserine Antibody, IgG	Less than 16 GPS
	Phosphatidylserine Antibody, IgM	Less than 22 MPS
GPS: IgG antiphosphatidylserine units, MPS: IgM antiphosphatidylserine units		

**HOTLINE NOTE:** There is a unit of measure change associated with this test.

Change the unit of measure for component 0050906, Phosphatidylserine Antibody IgG from U/mL to GPS.

Change the unit of measure for component 0050907, Phosphatidylserine Antibody IgM from U/mL to MPS.