

2011478

Arsenic, Random Urine with Reflex to Fractionated

U ARS RAND

**Specimen Required:** Patient Prep: Diet, medication, and nutritional supplements may introduce interfering substances. Patients should be encouraged to discontinue nutritional supplements, vitamins, minerals, nonessential over-the-counter medications (upon the advice of their physician), and avoid shellfish and seafood for 48 to 72 hours. High concentrations of iodine may interfere with elemental testing. **Collection of urine specimens from patients receiving iodinated or gadolinium-based contrast media should be avoided for a minimum of 72 hours post-exposure. Collection from patients with impaired kidney function should be avoided for a minimum of 14 days post contrast media exposure.**

Collect: Random urine.

Specimen Preparation: Transfer an 8 mL aliquot from a well-mixed collection to ARUP Trace Element-Free Transport Tubes (ARUP supply #43116), available online through eSupply using ARUP Connector contact ARUP Client Services at (800) 522-2787. (Min: 2 mL)

Storage/Transport Temperature: Refrigerated. Also acceptable: Room temperature or frozen.

Remarks: Trace Elements requisition form may be required (ARUP form #32990-Barcode; #32991-No Barcode).

Unacceptable Conditions: Acid preserved urine. **Specimens collected within 72 hours after administration of iodinated or gadolinium-based contrast media.**

Stability (collection to initiation of testing): Ambient: 1 week; Refrigerated: 2 weeks; Frozen: 1 year