

TEST CHANGE

Allergen, Food, Pineapple IgG 2011741, PNAPPL IGG	
Specimen Requirements:	
Patient Preparation:	
Collect:	<u>Plain red or serum</u> Serum separator tube (SST <u>)</u> .
Specimen Preparation:	Separate <u>serum</u> from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum or <u>plasma</u> to an ARUP <u>standard</u> <u>transport tube.Standard Transport Tube.</u> (Min: 0.252 mL). <u>Refer to "Allergen Specimen Collection Instructions" at</u> <u>www.aruplab.com/testing/resources/specimen.</u>)
Transport Temperature:	Refrigerated.
Unacceptable Conditions:	Hemolyzed, icteric, or lipemic specimens.
Remarks:	
Stability:	Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year
Methodology:	Quantitative ImmunoCAP Fluorescent Enzyme Immunoassay
Performed:	Sun
Reported:	1-8 days
Note:	
CPT Codes:	86001
New York DOH Approval Status:	Specimens from New York clients will be sent out to a New York DOH approved laboratory, if possible.
Interpretive Data:	
Values less than 2.00 mcg/mL represent absent or undetectable levels of allergen-specific IgG antibody.	
This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.	
Reference Interval:	
Less than 12.48 mcg/mL	

