

**TEST CHANGE** 

Phospholipase A2 Receptor (PLA2R) Antibody, IgG with Reflex to Titer 2011828, PLA2R

Specimen Requirements:

**Patient Preparation:** 

Collect: Serum Separator Tube

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection.

Transfer 1 mL serum to an ARUP Standard Transport Tube.

Effective Date: January 21, 2025

(Min: 0.2 mL)

Transport Temperature: Refrigerated

Unacceptable Conditions: Hemolyzed, hyperlipemic, icteric, heat-treated or contaminated

Remarks:

Stability: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1

month.1year

Methodology: Semi-Quantitative <u>Cell-Based</u> Indirect Fluorescent Antibody

Performed: Mon, Wed, Fri

Reported: 1-6 days

Note: If Phospholipase A2 Receptor Antibody, IgG is positive, then a

Phospholipase Receptor A2 Antibody, IgG titer will be added.

Additional charges apply.

CPT Codes: 86255; if reflexed, add 86256

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

A positive result (1:10 or greater) for phospholipase A2 receptor antibody, IgG in conjunction with other laboratory and clinical findings, supports a diagnosis of primary membranous glomerulonephritis (pMGN).

Reference Interval:

Less than 1:10