

TEST CHANGE

TEST CHANGE				
Smith and Smith/RNP (ENA) Antibodies, IgG				
3000460, SMI				
Specimen Requirements:				
Patient Preparation:				
Collect:		Serum Separator Tube (SST).		
Specimen Preparation:		Separate from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP Standard Transport Tube. (Min: 0.2 mL)		
Transport Temperature:		Refrigerated.		
Unacceptable Conditions:		Plasma or other body fluids. Contaminated, hemolyzed, or severely lipemic specimens.		
Remarks:				
Stability:		After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month (avoid repeated freeze/thaw cycles)		
Methodology:		Semi-Quantitative Enzyme-Linked Immunosorbent Assay/Semi-Quantitative Multiplex Bead Assay		
Performed:		Sun-Sat		
Reported:		1- <u>3</u> 2 days		
Note:				
CPT Codes:		86235 x2		
New York DOH Approval Status:		: This test is New York DOH approved.		
Interpretive Da	ita:			
Components	Interpretation			
Smith/RNP (ENA) Antibody, IgG	19 Units or less: Negative 20-39 Units: Weak Positive 40-80 Units: Moderate Positive 81 Units or greater: Strong Positive			
Smith (ENA) Antibody, IgG	29 AU/mL or less: Negative 30-40 AU/mL: Equivocal 41 AU/mL or greater: Positive			



Reference Interval:

Test Number	Components	Reference Interval
	Smith/RNP (ENA) Ab, IgG	19 Units or less
	Smith (ENA) Antibody, IgG	40 AU/mL or less