

TEST CHANGE

Smith and Smith/RNP (ENA) Antibodies, IgG

3000460, SMITH_RNP

Specimen Requirements:

Patient Preparation:

Collect: Serum separator tube (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP standard transport tube. (Min: 0.5 mL)

Transport Temperature: Refrigerated.

Unacceptable Conditions: Plasma or other body fluids. Contaminated, hemolyzed, or severely lipemic specimens.

Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 ~~year~~~~month~~ (avoid repeated freeze/thaw cycles)

Methodology: ~~Semi-Quantitative Enzyme-Linked Immunosorbent Assay (ELISA)~~ / Semi-Quantitative Multiplex Bead Assay

Note:

CPT Codes: 86235 x2

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Components	Interpretation
Smith/RNP (ENA) Ab Antibody, IgG	29 AU/mL 19 Units or less: Negative 30-20-39 Units: Weak Positive 40 AU/mL: Equivocal 41 AU/mL-80 Units: Moderate Positive 81 Units or greater: Strong Positive
Smith (ENA) Antibody, IgG	29 AU/mL or less: Negative 30-40 AU/mL: Equivocal 41 AU/mL or greater: Positive

Reference Interval:

Test Number	Components	Reference Interval
	Smith (ENA) Antibody, IgG	40 AU/mL or less
	Smith/RNP (ENA) Ab, IgG	40 AU/mL 19 Units or less

