

TEST CHANGE

Criteria Systemic Sclerosis Panel

3000479, SSC PANEL

Specimen Requirements:

Patient Preparation:

Collect: Serum ~~separator tube~~ **Separator Tube** (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 3 mL serum to an ARUP ~~standard transport tube~~ **Standard Transport Tube**. (Min: 0.25 mL)

Transport Temperature: Refrigerated.

Unacceptable Conditions: Contaminated, hemolyzed, or severely lipemic specimens.

Remarks:

Stability: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 ~~month~~ **year** (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative Indirect Fluorescent Antibody/Semi-Quantitative Multiplex Bead Assay/Semi-Quantitative Enzyme-Linked Immunosorbent Assay

Performed: Sun, Tue, Thu

Reported: 1-4 days

Note:

CPT Codes: 86039; 86235; 83516

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Component	Interpretation
Scleroderma (Scl-70) (ENA) Antibody, IgG	29 AU/mL or less Negative 30-40 AU/mL Equivocal 41 AU/mL or greater Positive
RNA Polymerase III Antibody, IgG	19 Units or less Negative 20-39 Units Weak Positive 40-80 Units Moderate Positive 81 Units or greater Strong Positive

Reference Interval:

Test Number	Components	Reference Interval
	Scleroderma (Scl-70) (ENA) Antibody, IgG	40 AU/mL or less
	RNA Polymerase III Antibody, IgG	19 Units or less
	Antinuclear Antibody (ANA), HEp-2, IgG	Less than 1:80