

Quarterly HOTLINE: Effective July 2, 2018

**New Test**  
Available Now

**3000485**

**Aldosterone Left Adrenal Vein**

**ALDO LAV**



Additional Technical Information

**Methodology:** Quantitative Chemiluminescent Immunoassay  
**Performed:** Sun-Sat  
**Reported:** Within 24 hours

**Specimen Required:** Collect: Adrenal venous sampling procedure is required. Serum Separator Tube (SST) or Plain Red.  
Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP Standard Transport Tube. (Min: 0.5 mL)  
Storage/Transport Temperature: Frozen.  
Unacceptable Conditions: EDTA plasma.  
Stability (collection to initiation of testing): After separation from cells: Ambient: 8 hours; Refrigerated: 5 days; Frozen: 1 month

**Note:** Refer to the Additional Technical Information link for information on Endocrine Society recommendations regarding patient preparation, specimen collection, medications for hypertension control during confirmatory testing for primary aldosteronism, and factors that may lead to false-positive or false-negative aldosterone-renin ratio (ARR) results.

**CPT Code(s):** 82088

New York DOH Approved.

**HOTLINE NOTE:** Refer to the Test Mix Addendum for interface build information.