

TEST CHANGE

Hepatitis C Virus (HCV) by Quantitative NAAT with Reflex to HCV High-Resolution Genotype by Sequencing 3000577, HCVQT HGR Specimen Requirements: Patient Preparation: Collect: Lavender (EDTA), pink (K2EDTA), plasma preparation tube (PPT), or serum separator tube (SST). Specimen Preparation: Separate from cells within 246 hours of collection. Transfer 3 mL serum or plasma to an ARUP standard transport tube (ARUP supply #15824). Available online through eSupply using ARUP Connect(TM) or contact ARUP Client Services at 800-522-2787. (Minimum volume. (Min: 1.87 mL) Transport Temperature: Frozen-Unacceptable Conditions: Heparinized specimens-Remarks: Stability: After separation from cells: Room temperature. Ambient: 24 hours (Critical: <u>s</u>hip FROZEN); Refrigerated: 72 hours; Frozen: 3 months6 weeks **Quantitative Polymerase Chain Reaction** Methodology: (PCR)/SequencingTranscription-Mediated Amplification (TMA) Performed: Sun-Sat Reported: 1-11 days Note: If Hepatitis C Virus by Quantitative NAAT result is greater than or equal to 16100,000 IU/mL, then Hepatitis C Virus High-Resolution Genotype by Sequencing will be added. Additional charges apply. CPT Codes: 87522; if reflexed, add 87902 New York DOH Approval Status: This test is New York DOH approved. Interpretive Data: Refer to report. **Beference Interval:**



Test Number	•	Reference Interval
	HCV Qnt by NAAT Interp	Not Detected

HOTLINE NOTE: There is a numeric map change associated with this test. Refer to the Hotline Test Mix for interface build information.