

Quarterly HOTLINE: Effective November 12, 2018

New Test	3000876	<i>Aspergillus fumigatus</i> Antibody IgG	ASPERF IGG
-----------------	----------------	--	-------------------

Methodology: Quantitative ImmunoCAP Fluorescent Enzyme Immunoassay

Performed: Sun

Reported: 1-8 days

Specimen Required: Collect: Serum Separator Tube (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum to an ARUP Standard Transport Tube. (Min: 0.2 mL)

Storage/Transport Temperature: Refrigerated.

Unacceptable Conditions: Hemolyzed, icteric, or lipemic specimens.

Stability (collection to initiation of testing): After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year

Reference Interval:

0.00-90.00 mcg/mL	Negative - No significant level of <i>A. fumigatus</i> IgG antibody detected.
90.01-99.99 mcg/mL	Equivocal - Questionable presence of <i>A. fumigatus</i> IgG antibody.
100.00 mcg/mL or greater	Positive – <i>A. fumigatus</i> IgG antibody detected. A positive result satisfies a single criterion in the determination of allergic bronchopulmonary aspergillosis (ABPA).

Interpretive Data: See Compliance Statement B: www.aruplab.com/CS

CPT Code(s): 86606

New York DOH Approved.

HOTLINE NOTE: Refer to the Test Mix Addendum for interface build information.