

TEST CHANGE

Hypersensitivity Pneumonitis 2

3001560, HYPER 2

Specimen Requirements:

Patient Preparation:

Collect: Serum ~~separator tube~~ **Separator Tube** (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 1 mL serum to an ARUP ~~standard transport tube~~ **Standard Transport Tube**. (Min: 0.15 mL)

Transport Temperature: Refrigerated.

Unacceptable Conditions: Plasma.

Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)

Methodology: Qualitative Immunodiffusion

Performed: ~~Sun-Sat~~ **Mon-Fri**

Reported: 3-~~7~~ **5** days

Note: Testing includes antibodies directed at Aspergillus fumigatus #2, A. fumigatus #3, A. flavus, Saccharomonospora viridis, and Thermoactinomyces candidus.

CPT Codes: 86331 x2; 86606 x3

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Reference Interval:

Test Number	Components	Reference Interval
	A. flavus Ab, Precipitin	None detected
	A. fumigatus #2 Ab, Precipitin	None detected
	A. fumigatus #3 Ab, Precipitin	None detected
	S. viridis Ab, Precipitin	None detected
	T. candidus Ab, Precipitin	None detected

