

HOTLINE: Effective August 19, 2019

New Test 3001782 Dermatomyositis Autoantibody Panel DERM PAN



Additional Technical Information

Methodology: Qualitative Immunoprecipitation/Qualitative Immunoblot

Performed: Sun-Sat **Reported:** 7-18 days

Specimen Required: Collect: Serum Separator Tube (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer one 1 mL serum aliquots to ARUP

Standard Transport Tubes. (Min: 0.5 mL/aliquot) Storage/Transport Temperature: Refrigerated.

<u>Unacceptable Conditions:</u> Hemolyzed, hyperlipemic, icteric, heat-treated or contaminated specimens. <u>Stability (collection to initiation of testing):</u> Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year

Reference Interval:

Test Number	Components	Reference Interval
	Mi-2 (nuclear helicase	Negative
	protein) Antibody	
	P155/140 Antibody	Negative
	SAE1 (SUMO activating	Negative
	enzyme) Ab	
	MDA5 (CADM-140) Ab	Negative
	NXP2 (Nuclear matrix	Negative
	protein-2) Ab	
	TIF1-gamma Ab	Negative

Interpretive Data: Refer to report.

See Compliance Statement D: www.aruplab.com/CS

Note: Antibodies: Mi-2, P155/140, SAE1, MDA5, NXP2, TIF1-gamma

CPT Code(s): 83516 x6

New York DOH Approved.

HOTLINE NOTE: Refer to the Test Mix Addendum for interface build information.