

**TEST CHANGE**

**Interstitial Lung Disease Autoantibody Panel**

3001784, ILD PANEL

**Specimen Requirements:**

**Patient Preparation:**

**Collect:** Serum Separator Tube (SST).

**Specimen Preparation:** Separate from cells ASAP or within 2 hours of collection. Transfer five 1 mL serum aliquots to ARUP Standard Transport Tubes. (Min: 2.8 mL)

**Transport Temperature:** Refrigerated.

**Unacceptable Conditions:** Hemolyzed, hyperlipemic, icteric, heat-treated or contaminated specimens.

**Remarks:**

**Stability:** Ambient: 24 hours; Refrigerated: 1 weeks; Frozen: 1 month

**Methodology:** Qualitative Immunoprecipitation/Semi-Quantitative Multiplex Bead Assay/Qualitative Immunoblot/Semi-Quantitative Enzyme-Linked Immunosorbent Assay/Quantitative Immunoturbidimetry

**Performed:** Sun-Sat

**Reported:** 7-18 days

**Note:** Antibodies: Ro52, Ro60, Jo-1, PL-7, PL12, EJ, Ku, SRP, OJ, PM/Scl-100, MDA5, CCP, Scl-70, RA, ANA, NXP-2, RNA Polymerase III

**CPT Codes:** 86235 x5; 83516 x7; 84182 x2; 86431; 86200; 86039

**New York DOH Approval Status:** This test is New York DOH approved.

**Interpretive Data:**

~~Refer to report.~~ ~~Refer to report.~~

~~This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.~~

Component	Interpretation	
SSA-52 (Ro52) (ENA) Antibody, IgG	29 AU/mL or Less 30-40 AU/mL 41 AU/mL or greater	Negative Equivocal Positive
SSA-60 (Ro60) (ENA) Antibody, IgG	29 AU/mL or Less 30-40 AU/mL 41 AU/mL or greater	Negative Equivocal Positive
Scleroderma (Scl-70) (ENA) Antibody, IgG	29 AU/mL or less 30-40 AU/mL 41 AU/mL or greater	Negative Equivocal Positive
Jo-1 Antibody, IgG	29 AU/mL or less 30-40 AU/mL 41 AU/mL or greater	Negative Equivocal Positive
Cyclic Citrullinated Peptide (CCP) Ab/Antibody, IgG/A	19 Units or less 20-39 Units 40-59 Units 60 Units or Greater	Negative Weak positive Moderate positive Strong positive
RNA Polymerase III Antibody, IgG	19 Units or less 20-39 Units 40-80 Units 81 Units or greater	Negative Weak positive Moderate positive Strong positive

Reference Interval:

Test Number	Components	Reference Interval
	Rheumatoid Factor	0-14 IU/mL
	Antinuclear Antibody (ANA), HEp-2, IgG	Less than 1:80
	Scleroderma (Scl-70) (ENA) Antibody, IgG	40 AU/mL or less
	SSA-52 (Ro52) (ENA) Antibody, IgG	40 AU/mL or less
	<del>Cyclic Citrullinated Peptide Ab, IgG</del>	<del>0-19 Units</del>
	Jo-1 (Histidyl-tRNA Synthetase) Ab, IgG	40 AU/mL or less
	PL-7 (threonyl-tRNA synthetase) Antibody	Negative
	PL-12 (alanyl-tRNA synthetase) Antibody	Negative
	EJ (glycyl-tRNA synthetase) Antibody	Negative
	Ku Antibody	Negative
	SRP (Signal Recognition Particle) Ab	Negative
	OJ (isoleucyl-tRNA synthetase) Antibody	Negative
	SSA-60 (Ro60) (ENA) Antibody, IgG	40 AU/mL or less
	MDA5 (CADM-140) Ab	Negative
	NXP2 (Nuclear matrix protein-2) Ab	Negative
	PM/Scl 100 Antibody, IgG	Negative

	RNA Polymerase III Antibody, IgG	19 Units or less
	<u>Cyclic Citrullinated Peptide Ab, IgG/A</u>	<u>19 Units or less</u>

---

**HOTLINE NOTE:** There is a component change associated with this test. One or more components have been added or removed. Refer to the Hotline Test Mix for interface build information.