

## TEST CHANGE

### Anti-sp100 and anti-gp210 Antibodies, IgG

3002482, SP100GP210

#### Specimen Requirements:

##### Patient Preparation:

Collect: Serum ~~separator tube~~ **Separator Tube** (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 1.0 mL serum to an ARUP ~~standard transport tube~~ **Standard Transport Tube**. (Min: 0.3 mL)

Transport Temperature: Refrigerated.

Unacceptable Conditions: ~~Nonserum~~ **Non-serum**, heat-inactivated, contaminated, grossly icteric, severely lipemic, grossly hemolyzed specimens, or inclusion of fibrin clot.

##### Remarks:

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative Enzyme-Linked Immunosorbent Assay

Performed: ~~Thu~~ **Wed**

Reported: 1-8 days

##### Note:

CPT Codes: 83516 x2

New York DOH Approval Status: This test is New York DOH approved.

#### Interpretive Data:

##### Refer to report

Component	Interpretation
Anti-sp100 Antibody, IgG	20.0 Units or less Negative 20.1-24.9 Units Equivocal 25.0 Units or greater Positive
Anti-gp210 Antibody, IgG	20.0 Units or less Negative 20.1-24.9 Units Equivocal 25.0 Units or greater

	Positive

Reference Interval:

Test Number	Components	Reference Interval
	Anti-gp210 Antibody, IgG	24.9 Units or less
	Anti-sp100 Antibody, IgG	24.9 Units or less