

TEST CHANGE

Interleukin 17, Plasma

3002626, IL17 PLA

Specimen Requirements:

Patient Preparation:

Collect: Green (~~lithium heparin~~ ~~Lithium Heparin~~).

Specimen Preparation: **CRITICAL FROZEN:** Separate plasma from cells ~~ASAP or~~ within 2 hours of collection. Transfer 1 mL plasma to an ARUP ~~standard transport tube.~~ ~~Standard Transport Tube.~~ (Min: 0.4 mL) **Freeze plasma within 30 minutes.**

Transport Temperature: CRITICAL FROZEN. Separate specimens must be submitted when multiple tests are ordered. Ship sample in an ARUP ~~standard transport tube.~~ ~~Standard Transport Tube.~~

Unacceptable Conditions: Contaminated or heat-inactivated specimens.

Remarks:

Stability: After separation from cells: Ambient: 30 minutes; Refrigerated: Unacceptable; Frozen: 1 year

Methodology: Quantitative Multiplex Bead Assay

Note: Cytokine levels may demonstrate diurnal variation. For longitudinal comparison, it is recommended that cytokine levels be determined at the same time of day.

CPT Codes: 83520

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Results are used to understand the pathophysiology of immune, infectious, or inflammatory disorders, or may be used for research purposes.

Reference Interval:

~~2.2 pg/mL or less~~

Test Number	Components	Reference Interval
	Interleukin 17, Plasma	2.2 pg/mL or less