
New Test	3002630	Interleukin 2, Plasma	IL2 PLA
-----------------	----------------	------------------------------	----------------

Methodology: Quantitative Multiplex Bead Assay
Performed: Sun-Sat
Reported: 1-4 days

Specimen Required: Collect: Green (lithium heparin).
Specimen Preparation: Separate plasma from cells ASAP or within 2 hours of collection. Transfer 1 mL plasma to an ARUP Standard Transport Tube. (Min: 0.4 mL)
Storage/Transport Temperature: **CRITICAL FROZEN. Separate specimens must be submitted when multiple tests are ordered. Ship in an ARUP Standard Transport Tube.**
Unacceptable Conditions: Refrigerated specimens. Contaminated or heat-inactivated specimens.
Stability (collection to initiation of testing): After separation from cells: Ambient: 30 minutes; Refrigerated: Unacceptable; Frozen: 1 year

Reference Interval:
2.1 pg/mL or less

Interpretive Data:
Results are used to understand the pathophysiology of immune, infectious, or inflammatory disorders, or may be used for research purposes.
See Compliance Statement B: www.aruplab.com/CS

Note: Cytokine levels may demonstrate diurnal variation. For longitudinal comparison, it is recommended that cytokine levels be determined at the same time of day.

CPT Code(s): 83520

New York DOH Approved.

HOTLINE NOTE: Refer to the Test Mix Addendum for interface build information.