

TEST CHANGE

Paraneoplastic Reflexive Panel 3002929, PNS PAN2		
Specimen Requirements:		
Patient Preparation:		
Collect:	Serum Separator Tube (SST)	
Specimen Preparation:	Separate from cells ASAP or within 2 hours of collection. Transfer 3 mL serum aliquot to an ARUP <u>standard transport</u> <u>tube.Standard Transport Tube.</u> (Min: 1.0 mL)	
Transport Temperature:	Refrigerated	
Unacceptable Conditions:	Contaminated, heat-inactivated, hemolyzed, or lipemic specimens	
Remarks:		
Stability:	After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month	
Methodology:	Semi-Quantitative Cell-Based Indirect Fluorescent Antibody/Qualitative Immunoblot	
Performed:	Wed	
Reported:	1-9 days	
Note:	Purkinje Cell (PCCA) antibody and Neuronal Nuclear (ANNA) antibody IgG are screened by IFA. If the IFA screen is indeterminate, then a Neuronal Nuclear Antibodies (Hu, Ri, Yo, Tr/DNER) IgG by Immunoblot will be added. If the IFA screen is positive at 1:10 or greater, then a PCCA/ANNA antibodies titer and Neuronal Nuclear Antibodies (Hu, Ri, Yo, Tr/DNER) IgG by Immunoblot will be added. Additional charges apply. If CV2.1 Antibody IgG Screen by IFA is positive, then CV2.1 Antibody IgG Titer by IFA will be added. Additional charges apply.	
CPT Codes:	86255 x2; 84182 x2; if reflexed add 86256 and/or 84182 x4; if reflexed add 86256	
New York DOH Approval Status:	This test is New York DOH approved.	
Interpretive Data:		
Refer to report		



Reference Interval:

Test Number	Components	Reference Interval
	Neuronal Antibody (Amphiphysin)	Negative
	Purkinje Cell/Neuronal Nuclear IgG Scrn	None Detected
	CV2-1 Ab IgG CBA-IFA Screen, Serum	Less than 1:10 <mark>0</mark>
	SOX1 Antibody, IgG by Immunoblot, Serum	Negative

HOTLINE NOTE: There is a reflexive pattern change associated with this test. One or more orderable or component has been added or removed to the reflexive pattern. Refer to the Hotline Test Mix for interface build information.