

HOTLINE: Effective October 20, 2020

New Test 3003255 Borrelia burgdorferi VlsE1/pepC10 Antibodies, Total by ELISA LPEP CHRON

with Reflex to IgG by Immunoblot

Available Now

Methodology: Semi-Quantitative Enzyme-Linked Immunosorbent Assay/Qualitative Immunoblot

Performed: Mon-Fri **Reported:** 1-4 days

Specimen Required: Collect: Serum separator tube.

Specimen Preparation: Separate serum from cells ASAP or within 2 hours of collection. Transfer 2 mL serum to an ARUP Standard

Transport Tube. (Min: 0.15 mL)

Storage/Transport Temperature: Refrigerated.

<u>Unacceptable Conditions:</u> Plasma, CSF. Contaminated, heat-inactivated, hemolyzed, icteric, or lipemic specimens.

Stability (collection to initiation of testing): After separation from cells: Ambient: 48 hours; Refrigerated: 10 days; Frozen: 1 month

Reference Interval:

Test Number	Components	Reference Interval	
3003253	Borrelia burgdorferi VlsE1/pepC10 Antibodies, Total by ELISA	0.90 IV or less	Negative – VlsE1 and pepC10 antibodies to <i>B. burgdorferi</i> not detected.
		0.91-1.09 IV	Equivocal – Repeat testing in 10-14 days may be helpful.
		1.10 IV or greater	Positive – VlsE1 and pepC10 antibodies to <i>B. burgdorferi</i> detected.
0050255	Borrelia burgdorferi Antibody, IgG by Immunoblot	Effective August 15, 2011 Negative	

Interpretive Data:

Refer to report

Note: If VIsE1/pepC10 antibodies by ELISA is 0.91 IV or greater, then IgG Immunoblot will be added. Additional charges apply.

CPT Code(s): 86618; if reflexed, add 86617

New York DOH Approved.

HOTLINE NOTE: Refer to the Test Mix Addendum for interface build information.