

## **TEST CHANGE**

## Vedolizumab Quantitation with Antibodies, Serum

3003676, VEDOL AB	
Specimen Requirements:	
Patient Preparation:	12 hours prior to specimen collection discontinue multivitamins or dietary supplements containing biotin (vitamin B7), commonly found in hair, skin, and nail supplements. Nivolumab (Opdivo) must be discontinued at least 4 weeks prior to testing.
Collect:	Plain red. Also acceptable: Serum separator tube (SST). Collect immediately before next scheduled dose (trough specimen).
Specimen Preparation:	Separate from cells ASAP or within 2 hours of collection. Transfer 1.5 mL serum to an ARUP standard transport tube. (Min: 0.75 mL) Test is not performed at ARUP; separate specimens must be submitted when multiple tests are ordered.
Transport Temperature:	Refrigerated. Also acceptable: Frozen.
Unacceptable Conditions:	
Remarks:	
Stability:	Ambient: Unacceptable; Refrigerated: 28 days; Frozen: 28 days
Methodology:	Quantitative Liquid Chromatography-Tandem Mass Spectrometry /Electrochemiluminescent Immunoassay (ECLIA)
Performed:	Varies
Reported:	<u>8-14</u> 5-11 days
Note:	
CPT Codes:	80280; 82397
New York DOH Approval Status:	This test is New York DOH approved.
Interpretive Data:	
Reference Interval:	
By Report	