

TEST CHANGE

Inflammatory Bowel Disease Differentiation Panel

3003748, IBD-PAN

Specimen Requirements:

Patient Preparation: N/A

Collect: Serum Separator Tube (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 1.5 mL serum to an ARUP [standard transport tube](#). ~~Standard Transport Tube~~. (Min: 0.6 mL)

Transport Temperature: Refrigerated.

Unacceptable Conditions: Contaminated, heat-inactivated, hemolyzed, or severely lipemic specimens.

Remarks: N/A

Stability: After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: [30 days](#)~~1 year~~ (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative Indirect Fluorescent Antibody (IFA)/Semi-Quantitative Enzyme Immunoassay (EIA)

Performed: Sun-Sat

Reported: 1-4 days

Note: This test may be a useful tool for distinguishing ulcerative colitis (UC) from Crohn disease (CD) in patients with suspected inflammatory bowel disease. ANCA IFA is simultaneously tested on ethanol- and formalin-fixed slides to allow differentiation of C- and P-ANCA patterns.

CPT Codes: 86036; 86671 x2

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Refer to report.

Reference Interval:

Test Number	Components	Reference Interval		
	S. cerevisiae Antibody, IgG			
		20.0 Units or less	Negative	
		20.1-24.9 Units	Equivocal	
		25.0 Units or greater	Positive	
	S. cerevisiae Antibody, IgA			
		20.0 Units or less	Negative	
		20.1-24.9 Units	Equivocal	
		25.0 Units or greater	Positive	
	ANCA IFA Titer	Less than 1:20		
	ANCA IFA Pattern	None Detected		