

NEW TEST

[Click for Pricing](#)

Autoimmune Movement Disorder Panel, Serum

3006206, AIMDS

Specimen Requirements:

Patient Preparation:

Collect: Serum separator tube (SST)

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer three 1 mL serum aliquots to ARUP standard transport tubes. (Min: 0.5 mL/aliquot)

Transport Temperature: Frozen

Unacceptable Conditions: Amniotic fluid, ocular fluid, peritoneal fluid, synovial fluid, CSF, or plasma. Contaminated, hemolyzed, icteric, or lipemic specimens.

Remarks:

Stability: After separation from cells: Ambient: 24 hours; Refrigerated: 1 week; Frozen: 30 days (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative Cell-Based Indirect Fluorescent Antibody/Semi-Quantitative Indirect Fluorescent Antibody (IFA)/Qualitative Immunoblot/Semi-Quantitative Enzyme-Linked Immunosorbent Assay (ELISA)/Quantitative Radioimmunoassay (RIA)

Performed: Varies

Reported: 3-10 days

Note: If NMDA antibody IgG is positive, then titer will be performed. Additional charges apply. If CV2.1 antibody IgG is positive, then titer will be added. Additional charges apply. PCCA/ANNA antibody IgG is screened by IFA. If the IFA screen is indeterminate, then a Neuronal Nuclear Antibodies (Hu, Ri, Yo, and Tr/DNER) IgG by Immunoblot will be performed. If the IFA screen is positive at 1:10 or greater, then a PCCA/ANNA antibodies titer and Neuronal Nuclear Antibodies (Hu, Ri, Yo, Tr/DNER) IgG by Immunoblot will be performed. Additional charges apply. If LGI1 antibody IgG is positive, then titer will be added. Additional charges apply. If CASPR2 antibody IgG is positive, then titer will be added. Additional charges apply. If DPPX antibody IgG by IFA is positive, then titer will be added.

Additional charges apply. If AMPA antibody IgG is positive, then titer will be added. Additional charges apply. If GABA-BR antibody IgG is positive, then titer will be added. Additional charges apply. If GABA-AR antibody IgG by IFA is positive, then titer will be added. Additional charges apply. If mGluR1 antibody IgG by IFA is positive, then titer will be added. Additional charges apply. If ITPR1 antibody IgG by IFA is positive, then titer will be added. Additional charges apply. If IgLON5 antibody IgG by IFA is positive, then titer will be added. Additional charges apply.

CPT Codes: 86341; 86596; 84182 x2; 86255 x12; if reflexed add 84182 x4; 86256 per titer

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Refer to report.

Component	Interpretation
P/Q-Type Voltage-Gated Calcium Channel (VGCC) Antibody	0.0-24.5 pmol/L Negative 24.6-45.6 pmol/L Indeterminate 45.7 pmol/L or greater Positive

Reference Interval:

Test Number	Components	Reference Interval
	CV2.1 Antibody IgG Screen by IFA	Less than 1:10
	Neuronal Antibody (Amphiphysin)	Negative
	Glutamic Acid Decarboxylase Antibody	0.0-5.0 IU/mL
	Purkinje Cell/Neuronal Nuclear IgG Scrn	None Detected
	N-methyl-D-Aspartate Receptor Ab, Serum	Less than 1:10
	P/Q-Type Calcium Channel Antibody	24.5 pmol/L or less
	SOX1 Antibody, IgG by Immunoblot, Serum	Negative
	AMPA Receptor Ab IgG Screen, Serum	Less than 1:10
	GABA-B Receptor Ab IgG Screen, Serum	Less than 1:10
	DPPX Ab IgG CBA IFA Screen, Serum	Less than 1:10
	GABA-AR Ab IgG CBA-IFA Screen, Serum	Less than 1:10
	IgLON5 Ab IgG CBA-IFA Screen, Serum	Less than 1:10
	mGluR1 Ab IgG CBA-IFA Screen, Serum	Less than 1:10
	ITPR1 Ab IgG CBA-IFA Screen, Serum	Less than 1:10
	CASPR2 Ab IgG Screen by IFA, Serum	Less than 1:10
	LG11 Ab IgG Screen by IFA, Serum	Less than 1:10

HOTLINE NOTE: Refer to the Hotline Test Mix for interface build information.