

NEW TEST

[Click for Pricing](#)

Autoimmune Stiff-Person Disorders, Serum

3006234, AISPSS

Specimen Requirements:

Patient Preparation: N/A

Collect: Serum separator tube (SST)

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer three 1 mL serum aliquots to ARUP standard transport tubes. (Min: 0.5 mL/aliquot)

Transport Temperature: Frozen

Unacceptable Conditions: Amniotic fluid, ocular fluid, peritoneal fluid, synovial fluid, CSF, or plasma. Contaminated, hemolyzed, icteric, or lipemic specimens.

Remarks:

Stability: After separation from cells: Ambient: 24 hours; Refrigerated: 1 week; Frozen: 30 days (avoid repeated freeze/thaw cycles)

Methodology: Semi-Quantitative Cell-Based Indirect Fluorescent Antibody/Qualitative Immunoblot/Semi-Quantitative Enzyme-Linked Immunosorbent Assay (ELISA)

Performed: Varies

Reported: 3-10 days

Note: If DPPX antibody IgG by IFA is positive, then titer will be added. Additional charges apply.

CPT Codes: 86341; 84182; 86255; if reflexed, add 86256

New York DOH Approval Status: This test is New York DOH approved.

Interpretive Data:

Refer to report.

Reference Interval:

Test Number	Components	Reference Interval
	Neuronal Antibody (Amphiphysin)	Negative
	Glutamic Acid Decarboxylase Antibody	0.0-5.0 IU/mL
	DPPX Ab IgG CBA IFA Screen, Serum	Less than 1:10

HOTLINE NOTE: Refer to the Hotline Test Mix for interface build information.