

## NEW TEST – Available Now

Click for Pricing

## Allergen, Food, Brazil Nut Component Ber e 1, IgE

3018631, BER E 1

Specimen Requirements:	
Patient Preparation:	Multiple patient encounters should be avoided.
Collect:	Serum separator tube.
Specimen Preparation:	Separate serum from cells ASAP or within 2 hours of collection. Transfer 0.5 mL serum to an ARUP standard transport tube. (Min: 0.25 mL). For multiple allergen orders refer to "Allergen Specimen Collection Instructions" at www.aruplab.com/testing/resources/specimen.
Transport Temperature:	Refrigerated.
Unacceptable Conditions:	Postmortem samples
Remarks:	
Stability:	After separation from cells: Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month
Methodology:	Quantitative ImmunoCAP Fluorescent Enzyme Immunoassay
Performed:	Sun-Sat
Reported:	1-3 days
Note:	
CPT Codes:	86008
New York DOH Approval Status:	This test is New York DOH approved.
Interpretive Data:	



Allergen results of 0.10-0.34 kU/L are intended for specialist use as the clinical relevance is undetermined. Even though increasing ranges are reflective of increasing concentrations of allergen-specific IgE, these concentrations may not correlate with the degree of clinical response or skin testing results when challenged with a specific allergen. The correlation of allergy laboratory results with clinical history and in vivo reactivity to specific allergens is essential. A negative test may not rule out clinical allergy or even anaphylaxis.

Reporting Range (reported in kU/L)	Probability of IgE Mediated Clinical Reaction	Class Scoring	
Less than 0.10	No significant level detected	0	
0.10-0.34	Clinical relevance undetermined	0/1	
0.35-0.70	Low	1	
0.71-3.50	Moderate	2	
3.51-17.50	High	3	
17.51-50.00	Very high	4	
20.01-100.00	Very high	5	
Greater than 100.00	Very high	6	

## Reference Interval:

Test Number	•	Reference Interval
	Brzl Nut Ber e1	Less than or equal to 0.09 kU/L

HOTLINE NOTE: Refer to the Hotline Test Mix for interface build information.