

**NEW TEST**

[Click for Pricing](#)

**Dermatomyositis and Polymyositis Panel**

3018866, COMBI PAN2

**Specimen Requirements:**

**Patient Preparation:**

**Collect:** Serum separator tube (SST).

**Specimen Preparation:** Separate from cells ASAP or within 2 hours of collection. Transfer two 1 mL serum aliquots to ARUP standard transport tubes. (Min: 0.5 mL/aliquot)

**Transport Temperature:** Refrigerated.

**Unacceptable Conditions:** Hemolyzed, hyperlipemic, icteric, heat-treated, or contaminated specimens

**Remarks:**

**Stability:** Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 month

**Methodology:** Qualitative Immunoprecipitation / Semi-Quantitative Multiplex Bead Assay / Qualitative Immunoblot / Semi-Quantitative Indirect Fluorescent Antibody (IFA)

**Performed:** Sun-Sat

**Reported:** 7-18 days

**Note:** Antibodies: PL-7, PL12, EJ, OJ, SRP, Jo-1, Mi-2, P155/140, SAE1, MDA5, NXP2, TIF1-gamma, ANA, Ha, Ks, Zo

**CPT Codes:** 83516 x7; 84182 x7; 86235; 86039

**New York DOH Approval Status:** This test is New York DOH approved.

**Interpretive Data:**

**Reference Interval:**

Test Number	Components	Reference Interval
-------------	------------	--------------------

**HOTLINE NOTE: Refer to the Hotline Test Mix for interface build information.**