

Periodic Fever Syndromes Panel, Sequencing and Deletion/Duplication

Last Literature Review: January 2022 Last Update: December 2023

Periodic fever syndromes are a varied group of autoinflammatory disorders characterized by recurrent episodes of fever that lack an infectious cause. These syndromes include familial Mediterranean fever (FMF), cyclic neutropenia, tumor necrosis factor receptor associated periodic syndrome (TRAPS), Muckle-Wells syndrome, and Hyper-IgD syndrome (HIDS). Genetic testing can confirm a diagnosis or be used to determine whether individuals with a family history of a periodic fever syndrome may be carriers.

Disease Overview

- For specific disease descriptions, refer to the [Genes Tested](#) table.
- Attacks often begin with a prodromal phase, with symptoms such as fatigue, malaise, and headache.
- Inflammatory symptoms (eg, fever, pain, rash) follow the prodromal phase.
- Symptoms usually resolve spontaneously.
- Individuals are generally asymptomatic between attacks, though in some severe cases, inflammatory symptoms may not completely resolve between attacks.
- Depending on the specific syndrome, symptoms may be triggered by exposure to cold or trauma.

Indications for Ordering

- Use to confirm diagnosis of a periodic fever syndrome in a symptomatic individual
- Diagnostic or carrier testing in individuals with a family history of a periodic fever syndrome

Prevalence

Varies by condition and ethnicity.

Inheritance

See the [Genes Tested](#) table.

Test Description

See the [Genes Tested](#) table for genes included in the panel.

Methodology

This test is performed using the following sequence of steps:

- Selected genomic regions, primarily coding exons and exon-intron boundaries, from the targeted genes are isolated from extracted genomic DNA using a probe-based hybrid capture enrichment workflow.
- Enriched DNA is sequenced by massively parallel sequencing (MPS), also known as next generation sequencing (NGS), followed by paired-end read alignment and variant calling using a custom bioinformatics pipeline. The pipeline includes an algorithm for detection of large (single exon-level or larger) deletions and duplications.
- Sanger sequencing is performed as necessary to fill in regions of low coverage and, in certain situations, to confirm variant calls.
- Large deletion/duplication calls made using MPS are confirmed by an orthogonal exon-level microarray when sample quality and technical conditions allow.

Clinical Sensitivity

Variable, dependent on specific disorder

Featured ARUP Testing

[Periodic Fever Syndromes Panel, Sequencing and Deletion/Duplication 2007370](#)

Method: Massively Parallel Sequencing

- Preferred test to confirm a diagnosis of a periodic fever syndrome
- Predictive diagnostic or carrier testing in individuals with a family history of a periodic fever syndrome

If a familial sequence variant has been previously identified, targeted sequencing for that variant may be appropriate; refer to the [Laboratory Test Directory](#) for additional information.

Analytic Sensitivity

Variant Class	Analytic Sensitivity (PPA) Estimate ^a (%) and 95% Credibility Region (%)	Analytic Specificity (NPA) (%)
SNVs	>99 (96.9-99.4)	>99.9
Deletions 1-10 bp ^b	93.8 (84.3-98.2)	>99.9
Insertions 1-10 bp ^b	94.8 (86.8-98.5)	>99.9
Exon-level ^c Deletions	97.8 (90.3-99.8) [2 exons or larger] 62.5 (38.3-82.6) [single exon]	>99.9
Exon-level ^c Duplications	83.3 (56.4-96.4) [3 exons or larger]	>99.9

^aGenes included on this test are a subset of a larger methods-based validation from which the PPA values are derived. These values do not apply to testing performed by multiplex ligation-dependent probe amplification (MLPA).

^bVariants greater than 10 bp may be detected, but the analytical sensitivity may be reduced.

^cIn most cases, a single exon deletion or duplication is less than 450 bp and 3 exons span a genomic region larger than 700 bp.

bp, base pairs; NPA, negative percent agreement; PPA, positive percent agreement; SNVs, single nucleotide variants

Limitations

- A negative result does not exclude a heritable form of a periodic fever syndrome.
- Diagnostic errors can occur due to rare sequence variations.
- Interpretation of this test result may be impacted if the individual has had an allogeneic stem cell transplantation.
- The following will not be evaluated:
 - Variants outside the coding regions and intron-exon boundaries of the targeted genes
 - Regulatory region and deep intronic variants
 - Breakpoints of large deletions/duplications
- The following may not be detected:
 - Deletions/duplications/insertions of any size by massively parallel sequencing
 - Large duplications less than three exons in size
 - Noncoding transcripts
 - Low-level somatic variants
 - Certain other variants due to technical limitations in the presence of pseudogenes or repetitive/homologous regions

Genes Tested

Gene	MIM Number	Disorder	Inheritance	Age of Onset	Clinical Symptoms
<i>ELANE</i>	130130	Cyclic neutropenia	AD	Infancy	Fever and malaise Mouth ulcers Cyclic neutropenia Chronic and severe infections
		Severe congenital neutropenia 1	AD	Infancy	Fever Inflammation of gums and skin Decreased levels of neutrophils Chronic and severe infections
<i>LPIN2</i>	605519	Majeed syndrome	AR	Before 2 years	Recurrent fever episodes Chronic recurrent multifocal osteomyelitis (CRMO)

AD, autosomal dominant; AR, autosomal recessive

Gene	MIM Number	Disorder	Inheritance	Age of Onset	Clinical Symptoms
					<p>Congenital dyserythropoietic anemia</p> <p>Sweet syndrome—painful bumps and blisters</p> <p>Contractures</p> <p>Growth retardation</p>
<i>MEFV</i>	608107	Familial Mediterranean fever (FMF)	AR and AD	Typically before 10 years for AR FMF Milder/AD forms may not present until adulthood	<p>Recurrent fever episodes</p> <p>Painful inflammation in abdomen, chest, and joints</p> <p>Erysipelas-like rash on lower legs</p>
	608068	Acute febrile neutrophilic dermatosis	AD	Childhood	<p>Recurrent fever</p> <p>Acute onset of painful skin lesions</p> <p>Arthralgia and myalgia</p>
<i>MVK</i>	251170	Porokeratosis 3	AD	Between 3rd-4th decade of life	<p>Annular skin plaques surrounded by distinctive keratotic rim</p> <p>Fluctuate seasonally</p>
		Hyper-IgD syndrome (HIDS)	AR	Infancy	<p>Periodic high fevers</p> <p>Abdominal and joint pain</p> <p>Headache</p> <p>Skin lesions</p> <p>Hepatomegaly and/or splenomegaly</p> <p>Elevated immunoglobulin D</p>
		Mevalonic aciduria	AR	Infancy	<p>Hepatosplenomegaly</p> <p>Abdominal and joint pain</p> <p>Skin rashes</p> <p>Failure to thrive</p> <p>Developmental delay and progressive ataxia</p> <p>Progressive vision problems</p>
<i>NLRP12</i>	609648	Familial cold autoinflammatory syndrome 2 (FCAS2)	AD	First year of life to middle age	<p>Episodic and recurrent rash</p> <p>Urticaria</p> <p>Arthralgia</p> <p>Myalgia</p> <p>Abdominal and/or thoracic pain</p>
<i>NLRP3</i>	606416	Familial cold autoinflammatory syndrome 1 (FCAS1)	AD	Before age 10	<p>Recurrent episodes of nonpruritic urticaria rash</p> <p>Episodes triggered by exposure to cold</p> <p>Low-grade fever and malaise</p> <p>Sweating, headaches, and nausea</p>
		Keratoendotheliitis fugax hereditaria	AD	Between ages 4-12 years	<p>Periodic inflammation of corneal endothelium</p> <p>Redness of the eye, pain, and photophobia</p> <p>Blurry vision</p>

dominant; AR, autosomal recessive

Gene	MIM Number	Disorder	Inheritance	Age of Onset	Clinical Symptoms
		Muckle-Wells syndrome	AD	Infancy to early childhood	Recurrent rashes Intermittent fevers Joint pain Recurrent conjunctivitis Progressive hearing loss Amyloidosis
		Neonatal onset multisystem inflammatory disease (NOMID)/CINCA syndrome	AD	Infancy	Skin rash typically present at birth Chronic meningitis Headaches, seizures, and vomiting Intellectual disability and developmental delay Hearing and vision loss Joint inflammation and cartilage overgrowth Short stature Contractures
		Deafness, autosomal dominant 34, with or without inflammation	AD	Childhood	Postlingual sensorineural hearing loss Episodic urticaria Periodic fever Renal amyloidosis
NOD2	605956	Blau syndrome	AD	Early childhood; usually before age 4	Granulomatous dermatitis Arthritis Uveitis Nephritis; chronic kidney failure
PSTPIP1	606347	Pyogenic sterile arthritis, pyoderma gangrenosum, and acne	AD	Childhood	Pyogenic arthritis Pyoderma gangrenosum Severe cystic acne
TNFAIP3	191163	Autoinflammatory syndrome, familial, Behcet-like	AD	First or second decade of life	Mucosal ulcers (particularly in oral and genital areas) Skin rash Uveitis Polyarthritis
TNFRSF1A	191190	Tumor necrosis factor receptor-associated periodic syndrome (TRAPS) (aka periodic fever, familial)	AD	Childhood	Recurrent fever Sterile peritonitis/pleuritis Abdominal pain Myalgia Leukocytosis Elevated erythrocyte sedimentation rate

AD, autosomal dominant; AR, autosomal recessive

