

Calprotectin, Fecal by Immunoassay

Inflammatory bowel disease (IBD) represents a spectrum of chronic disorders that affect the gastrointestinal (GI) tract. Crohn disease (CD) and ulcerative colitis (UC) are the major IBD disorders.¹ Fecal calprotectin is a marker of gut inflammation with good sensitivity for detecting IBD.¹ Fecal lactoferrin, an iron-binding protein, is another useful marker of intestinal inflammation in IBD,² but more clinical evidence is available for fecal calprotectin.³

Disease Overview

Prevalence

IBD: 286/100,000 in United States⁴

Physiology

- Calprotectin is a calcium-binding protein² and accounts for 60% of soluble protein in neutrophils³
- Calprotectin concentration in feces is proportional to the level of inflammation in patients with UC³; the relationship is more variable in patients with CD⁵
- Calprotectin is stable in stool samples³

Diagnostic Issues

- IBD symptoms may be vague and similar to those of irritable bowel syndrome (IBS) (eg, diarrhea, abdominal pain)
 - IBS is much more prevalent than IBD
- Differentiation of IBD from IBS may require invasive procedures
- Calprotectin testing may be useful as a screen for differentiating IBS and IBD,² reducing the necessity of invasive procedures

Monitoring Issues

- Monitoring by endoscopy is invasive
- Calprotectin measurement can be used to help differentiate quiescent from active IBD³
- Mucosal healing is associated with sustained remission and is the goal of IBD treatment²
- Calprotectin levels correlate with endoscopic scoring systems that are used to assess mucosal healing and may be useful in evaluating mucosal healing³

Test Interpretation

Clinical Validation

Screening performance for IBD

- Sensitivity: 93% in adults; 92% in children⁶
- Specificity: 96% in adults; 76% in children⁶
- More sensitive and specific than serum inflammatory markers^{3,7}
- Individuals with high pretest probability of IBD (>75%) should be referred directly to endoscopy due to the risk of false-negative

Tests to Consider

Calprotectin, Fecal by Immunoassay 3002859

Method: Quantitative Chemiluminescent Immunoassay

- Aids in differentiation of IBD from IBS and other functional disorders of the GI system
 - Not specific for IBD
- Aids in monitoring IBD and prediction of relapse

See [Related Tests](#)

calprotectin results^{6,8}

- Screening for elevated fecal calprotectin in individuals with low pretest probability for IBD may result in cost savings by reducing need for unnecessary procedures⁸
 - Confirm positive results by endoscopy and follow negative result clinically



Results

Results	Range	Clinical Significance
Normal	≤50 µg/g	Likely to rule out IBD in adults with <75% prior probability
Borderline	51-120 µg/g	Reevaluation in 4-6 weeks is recommended
Abnormal	≥121 µg/g	Supports diagnosis of IBD

Limitations

- Calprotectin is not specific for IBD and is also elevated in:
 - GI infections
 - Colorectal cancer
 - Celiac disease
 - Mild elevations may be seen with nonsteroidal anti-inflammatory drug or aspirin use
- Calprotectin concentration alone is not diagnostic for IBD
- Calprotectin does not distinguish celiac disease from UC
 - Results may fluctuate as disease activity fluctuates
 - GI bleeding can cause mild increases in fecal calprotectin concentrations
- Concentrations of fecal biomarkers may vary in different stool samples from a single patient⁹

References

1. Yu YR, Rodriguez R. [Clinical presentation of Crohn's, ulcerative colitis, and indeterminate colitis: symptoms, extraintestinal manifestations, and disease phenotypes](#). *Semin Pediatr Surg.* 2017; 26 (6): 349-355. PubMed
2. Lichtenstein GR, Loftus EV, Isaacs KL, et al. [ACG clinical guideline: management of Crohn's disease in adults](#). *Am J Gastroenterol.* 2018; 113 (4): 481-517. PubMed
3. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. [ACG clinical guideline: ulcerative colitis in adults](#). *Am J Gastroenterol.* 2019; 114 (3): 384-413. PubMed
4. Georgy M, Negm Y, El-Matary W. [Quality improvement in healthcare for patients with inflammatory bowel disease](#). *Transl Pediatr.* 2019; 8 (1): 77-82. PubMed
5. Costa F, Mumolo MG, Ceccarelli L, et al. [Calprotectin is a stronger predictive marker of relapse in ulcerative colitis than in Crohn's disease](#). *Gut.* 2005; 54 (3): 364-8. PubMed
6. van Rheenen PF, Van de Vijver E, Fidler V. [Faecal calprotectin for screening of patients with suspected inflammatory bowel disease: diagnostic meta-analysis](#). *BMJ.* 2010;341:c3369. PubMed
7. Levine A, Koletzko S, Turner D, et al. [ESPGHAN revised porto criteria for the diagnosis of inflammatory bowel disease in children and adolescents](#). *J Pediatr Gastroenterol Nutr.* 2014; 58 (6): 795-806. PubMed
8. Yang Z, Clark N, Park KT. [Effectiveness and cost-effectiveness of measuring fecal calprotectin in diagnosis of inflammatory bowel disease in adults and children](#). *Clin Gastroenterol Hepatol.* 2014; 12 (2): 253-62.e2. PubMed
9. Calafat M, Cabré E, Mañosa M, et al. [High within-day variability of fecal calprotectin levels in patients with active ulcerative colitis: what is the best timing for stool sampling?](#) *Inflamm Bowel Dis.* 2015; 21 (5): 1072-6. PubMed

Related Information

[Inflammatory Bowel Disease - IBD](#)

Related Tests



[Lactoferrin, Fecal by ELISA 0061164](#)

Method: Qualitative Enzyme-Linked Immunosorbent Assay

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology. 500 Chipeta Way, Salt Lake City, UT 84108
(800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com
Content Review August 2020 | Last Update August 2020

© 2021 ARUP Laboratories. All Rights Reserved.

Client Services - (800) 522-2787