

Respiratory Pathogen Molecular Panel Testing

Last Literature Review: May 2019 Last Update: December 2023

Viral respiratory tract infections are the most common diseases affecting humans worldwide. Respiratory viruses can be associated with both self-limiting upper respiratory tract infections (eg, the common cold) and more severe lower respiratory tract infections (LRTIs) (eg, bronchitis, bronchiolitis, pneumonitis, pneumonia). LRTIs are a major cause of hospitalization, morbidity, and mortality in infants and the elderly and are associated with significant disease burden.

Disease Overview

Incidence

- Viral infections cause approximately 80% of respiratory tract disease; mixed infections account for approximately 20% infection in adults, 60% in children
- · Symptomatic disease varies by viral type and patient age
 - Highest rates occur in young and elderly populations with widely varying outcomes
- · Seasonal influenza causes a large proportion of viral respiratory disease
 - o Median incidence of approximately 8% in U.S. 2010 to 2016

Symptoms

Symptoms general occur within 1-3 days of exposure and last 7-14 days.

Location	Symptoms
Upper respiratory infections	Nasal congestion
	Sneezing
	Cough
	Sore throat
	Fever
	Chills
	Fatigue
	Decreased appetite ^a
	Lethargy ^a
Lower respiratory infections ^b	Worsening cough
	Shortness of breath
	Focal pain
	Dizziness ^c
	Confusion ^C
^a Especially in children	
^b Overlap with upper infection	
^c Older adults	

Featured ARUP Testing

Respiratory Virus Mini Panel by PCR 0060764

Method: Qualitative Reverse Transcription Polymerase Chain Reaction

Preferred test to confirm respiratory syncytial virus (RSV), influenza A, or influenza B in general inpatients and RSV in adults.

Respiratory Viral Panel by PCR 3001479

Method: Qualitative Polymerase Chain Reaction

- Preferred test for evaluating severely immunocompromised (eg, BMT) or critically ill (ICU) patients with respiratory symptoms.
- Test detects influenza A, influenza B, RSV, human metapneumovirus, human rhinovirus, and adapovirus
- Detects and differentiates parainfluenza 1, 2, 3, and 4.

Test Interpretation

Limitations

Negative result

- Does not rule out the presence of polymerase chain reaction (PCR) inhibitors in patient specimen
- Does not detect assay-specific nucleic acid in concentrations below level of detection by assay

Related Information

Respiratory Syncytial Virus - RSV Respiratory Viruses

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology. 500 Chipeta Way, Salt Lake City, UT 84108 (800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com

© 2024 ARUP Laboratories. All Rights Reserved.

Client Services - (800) 522-2787