

## Warfarin Sensitivity (CYP2C8, CYP2C9, CYP4F2, VKORC1) Genotyping

Warfarin (Coumadin) is an anticoagulant widely used throughout the world. Testing may be indicated for warfarin-naïve individuals prior to starting warfarin therapy, individuals with a personal or family history of difficulty with warfarin, and adherent patients taking warfarin but who are difficult to treat. Testing is also indicated for individuals currently on warfarin and required to discontinue therapy, such as prior to an invasive procedure or surgery, to estimate the time required to eliminate the drug.

### DISEASE OVERVIEW

#### Pathophysiology

- Warfarin is administered as a racemic mixture; s-warfarin is more potent than r-warfarin and is thought to mediate most of the anticoagulant activity of warfarin.
- Primary mechanism of action is to inhibit vitamin K epoxide reductase (VKOR)
  - VKOR recycles vitamin K and activates clotting factors II, VII, IX, and X
- Exerts anticoagulant effects by reducing the concentration of these activated clotting factors

#### Diagnostic Issues

- Individual response to warfarin varies:
  - Factors affecting response include age, gender, body mass, diet, concomitant medications, and genetic variants
  - An estimated 40-63% of the variability in therapeutic warfarin dose is accounted for by the *CYP2C9*\*2 and \*3 and the *VKORC1*\*2 variant alleles
- Overdosing and underdosing can result in life-threatening events (eg, bleeding or thrombosis).
  - ~1% of individuals die due to bleeding complications associated with warfarin.
  - ~15% of individuals experience minor bleeding complications.
- Dose adjustments are often necessary
  - Usually based on international normalized ratio (INR)
  - May be difficult to achieve therapeutic INR in some individuals (eg, those requiring <21 mg per week or >49 mg per week to maintain INR).

### Genetics

#### Genes Tested

*CYP2C8, CYP2C9, CYP4F2, VKORC1*

#### Variants Tested

Gene (Transcript)	Allele
<i>CYP2C8</i> (NM_000770)	<i>CYP2C8</i> *1C: rs17110453, c.-370T>G
	<i>CYP2C8</i> *2: rs11572103, c.805A>T
	<i>CYP2C8</i> *3: rs10509681, c.1196A>G

#### Tests to Consider

##### Warfarin Sensitivity (CYP2C8, CYP2C9, CYP4F2, VKORC1) Genotyping 3001541

**Method:** Polymerase Chain Reaction/Fluorescence Monitoring

Identifies individuals with inherited variants that affect metabolism (*CYP2C8, CYP2C9*) and/or sensitivity (*CYP4F2, VKORC1*) to warfarin.

See [Variants Tested](#) table for more information.

##### Related Test

##### CYP2C8 and CYP2C9 3001501

**Method:** Polymerase Chain Reaction/Fluorescence Monitoring

Assesses genetic risk of abnormal drug metabolism for *CYP2C8* and/or *CYP2C9* substrates.

Gene (Transcript)	Allele
	<i>CYP2C8*4</i> : rs1058930, c.792C>G
<i>CYP2C9</i> (NM_000771)	<i>CYP2C9*2</i> : rs1799853, c.430C>T
	<i>CYP2C9*3</i> : rs1057910, c.1075A>C
	<i>CYP2C9*4</i> : rs56165452, c.1076T>C
	<i>CYP2C9*5</i> : rs28371686, c.1080C>G
	<i>CYP2C9*6</i> : rs9332131, c.818del
	<i>CYP2C9*8</i> : rs7900194, c.449G>A
	<i>CYP2C9*11</i> : rs28371685, c.1003C>T
<i>CYP4F2</i> (NM_001082)	<i>CYP4F2*3</i> : rs2108622, c.1297G>A
<i>VKORC1</i> (NM_024006)	<i>VKORC1*2</i> : rs9923231, c.-1639G>A

Allele frequencies and phenotype predictions available at [www.pharmvar.org](http://www.pharmvar.org) or [www.pharmgkb.org](http://www.pharmgkb.org)

## Inheritance

Autosomal codominant

## Test Interpretation

### Sensitivity/Specificity

- Clinical sensitivity: genetic factors and known non-genetic factors account for ~50% of the variability in warfarin dose<sup>1</sup>
- Analytical sensitivity and specificity: >99%

### Results

- Variant(s) detected:
  - *VKORC1\*2* allele is associated with reduced expression of the warfarin target, vitamin K epoxide reductase (VKOR), and a reduced dose requirement
  - The *CYP4F2\*3* allele is associated with an increased dose requirement
  - *CYP2C8/9* variants are associated with a reduced rate of warfarin catabolism, which is associated with a decreased dose requirement and an increased time required to achieve steady state
    - Loading doses may be required
    - Vulnerability to drug-drug interactions may also be increased
- No variants detected: predictive of \*1 functional alleles
- Genotype should be interpreted with clinical information.
- Genetic information and nongenetic factors can be used in combination with warfarin dose calculators, such as through [www.WarfarinDosing.org](http://www.WarfarinDosing.org).
- Additional dosing guidance is available through drug labeling and professional guidance documents, such as those published by the American College of Chest Physicians (CHEST) and the Clinical Pharmacogenetics Implementation Consortium (CPIC).

### Limitations

- Only the targeted *CYP2C8*, *CYP2C9*, *CYP4F2* and *VKORC1* variants will be detected by this panel, and assumptions about phase and content are made to assign alleles.
- Diagnostic errors can occur due to rare sequence variations.
- Risk of therapeutic failure or adverse reactions with CYP2C8 or CYP2C9 substrates may be affected by genetic and nongenetic factors that are not detected by this test.
- This result does not replace the need for therapeutic drug or clinical monitoring.

## References

1. Clinical Pharmacogenetics Implementation Consortium. [CPIC guideline for pharmacogenetics-guided warfarin dosing](#). [Last modified: Apr 2019; Accessed: Apr 2019]

## Additional Resources

Kearon C, Akl EA, Ornelas J, et al. [Antithrombotic therapy for VTE disease: CHEST Guideline and Expert Panel Report](#). *Chest*. 2016;149(2):315-352. PubMed

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology, 500 Chipeta Way, Salt Lake City, UT 84108  
(800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com  
Content Review July 2019 | Last Update July 2020